



Available online at www.sciencedirect.com



Journal of Differential Equations

Journal of Differential Equations 421 (2025) 1-49

www.elsevier.com/locate/jde

Existence and concentration of normalized solutions for p-Laplacian equations with logarithmic nonlinearity $\stackrel{\text{\tiny{}^{\diamond}}}{=}$

Liejun Shen^a, Marco Squassina^{b,*}

^a Department of Mathematics, Zhejiang Normal University, Jinhua, Zhejiang, 321004, People's Republic of China ^b Dipartimento di Matematica e Fisica Università Cattolica del Sacro Cuore, Via della Garzetta 48, 25133, Brescia, Italy

Received 14 March 2024; revised 25 September 2024; accepted 26 November 2024

Abstract

We investigate the existence and concentration of normalized solutions for a p-Laplacian problem with logarithmic nonlinearity of type

$$\begin{cases} -\varepsilon^p \Delta_p u + V(x)|u|^{p-2}u = \lambda |u|^{p-2}u + |u|^{p-2}u \log |u|^p \text{ in } \mathbb{R}^N, \\ \int |u|^p dx = a^p \varepsilon^N, \\ \mathbb{R}^N \end{cases}$$

where $a, \varepsilon > 0, \lambda \in \mathbb{R}$ is known as the Lagrange multiplier, $\Delta_p \cdot = \operatorname{div}(|\nabla \cdot |^{p-2}\nabla \cdot)$ denotes the usual *p*-Laplacian operator with $2 \le p < N$ and $V \in C^0(\mathbb{R}^N)$ is the potential which satisfies some suitable assumptions. We prove that the number of positive solutions depends on the profile of *V* and each solution concentrates around its corresponding global minimum point of *V* in the semiclassical limit when $\varepsilon \to 0^+$ using variational method. Moreover, we also get the existence of normalized solutions for some logarithmic *p*-Laplacian equations involving mass-supercritical nonlinearities.

© 2024 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

MSC: 35J10; 35J20; 35B06

* Corresponding author. E-mail addresses: ljshen@zjnu.edu.cn (L. Shen), marco.squassina@unicatt.it (M. Squassina).

https://doi.org/10.1016/j.jde.2024.11.049

^{*} L.J. Shen is partially supported by NSFC (12201565). M. Squassina is member of Gruppo Nazionale per l'Analisi Matematica, la Probabilita e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

^{0022-0396/© 2024} The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

Keywords: Normalized solutions; Logarithmic *p*-Laplacian equation; Multiplicity; Singularly perturbed; Minimization technique; Variational method

Contents

1.	Introduction and main results	2
2.	Variational setting and preliminaries 1	1
3.	The semiclassical problem	9
4.	The autonomous problem	6
	4.1. The L^p -subcritical case	6
	4.2. The L^p -supercritical case	0
5.	Final comments	1
Ackno	owledgments	2
Appei	ndix A. Some technical stuff	2
Data a	availability	7
Refere	ences	7

1. Introduction and main results

In this article, we aim to establish the existence and concentrating behavior of nontrivial solutions for the following p-Laplacian equations with logarithmic nonlinearity of type

$$-\varepsilon^{p}\Delta_{p}u + V(x)|u|^{p-2}u = \lambda|u|^{p-2}u + |u|^{p-2}u\log|u|^{p} \text{ in } \mathbb{R}^{N},$$
(1.1)

under the constraint

$$\int_{\mathbb{R}^N} |u|^p dx = a^p \varepsilon^N,\tag{1.2}$$

where $a, \varepsilon > 0, \lambda \in \mathbb{R}$ is known as the Lagrange multiplier, $\Delta_p \cdot = \operatorname{div}(|\nabla \cdot|^{p-2}\nabla \cdot)$ denotes the usual *p*-Laplacian operator with $2 \le p < N$ and $V \in \mathcal{C}^0(\mathbb{R}^N)$ is the potential which satisfies

 $(V_1) \quad V \in \mathcal{C}^0(\mathbb{R}^N) \text{ and } -\infty < V_0 \triangleq \inf_{x \in \mathbb{R}^N} V(x) < V_\infty \triangleq \lim_{|x| \to +\infty} V(x) < +\infty;$ $(V_2) \quad V^{-1}(\{V_0\}) = \{x^1, x^2, \cdots, x^l\} \text{ with } x^1 = 0 \text{ and } x^i \neq x^j \text{ if } i \neq j \text{ for all } i, j \in \{1, 2, \cdots, l\}.$

In the case p = 2, Eq. (1.1) also comes from the study of solitary waves for the following nonlinear Schrödinger equation

$$i\frac{\partial\psi}{\partial t} + \Delta\psi - V(x)\psi + h(|\psi|^2)\psi = 0 \text{ in } (0,\infty) \times \mathbb{R}^N, \qquad (1.3)$$

where i is the imaginary unit, $\psi : (0, \infty) \times \mathbb{R}^N \to \mathbb{C}$, $V : \mathbb{R}^N \to \mathbb{R}$ is the potential, $h(e^{i\theta}z) = e^{i\theta}h(z)$ for $z \in \mathbb{C}$ and $\theta \in \mathbb{R}$. It is not difficult to see that any solution ψ of Eq. (1.3) with the Cauchy initial function $\psi(0, x)$ preserves the L^2 -mass, namely

$$\int_{\mathbb{R}^N} |\psi(t,x)|^2 \mathrm{d}x = \int_{\mathbb{R}^N} |\psi(0,x)|^2 \mathrm{d}x, \ \forall t \in (0,\infty).$$

Actually, Eq. (1.3) is usually adopted in the study of nonlinear optics and Bose-Einstein condensates, where ψ describes the state of the condensate and the L^2 -mass is the total number of atoms, see e.g. [31,51,60]. One significant motivation associated with Eq. (1.3) is the search for its standing wave solutions. The standing wave is a solution of the form $\psi(t, x) = e^{-i\lambda t}u(x)$, where $\lambda \in \mathbb{R}$ and $u : \mathbb{R}^N \to \mathbb{R}$ is a time-independent function that satisfies the nonlinear elliptic equation

$$-\Delta u + V(x)u = \lambda u + g(u) \text{ in } \mathbb{R}^N, \qquad (1.4)$$

with $g(u) = h(|u|^2)u$.

Usually, there are two directions to the study of standing waves of the Schrödinger equation (1.4). On the one hand, one can choose the frequency λ to be fixed and consider the existence of nontrivial solutions for it by investigating critical points of the variational functional \mathcal{J}_{λ} : $H^1(\mathbb{R}^N) \to \mathbb{R}$ defined by

$$\mathcal{J}_{\lambda}(u) = \frac{1}{2} \int_{\mathbb{R}^N} \left(|\nabla u|^2 + (V(x) - \lambda)|u|^2 \right) dx - \int_{\mathbb{R}^N} G(u) dx,$$

where $G(t) = \int_0^t g(s)ds$. When $g(t) = t \log t^2$, we refer to the article [62] and its references therein to acquaint the significant physical applications in quantum mechanics, quantum optics, nuclear physics, transport and diffusion phenomena, open quantum systems, effective quantum gravity, superfluidity theory and Bose-Einstein condensation for Eq. (1.4). Owing to the logarithmic type nonlinearity, it presents some challenging mathematical problems. For instance, the associated variational functional is not C^1 -smooth since one can find a function below

$$u(x) = \begin{cases} (|x|^{N/2}\log(|x|))^{-1}, & |x| \ge 3, \\ 0, & |x| \le 2, \end{cases}$$

such that $u \in H^1(\mathbb{R}^N)$, however it holds that $\int_{\mathbb{R}^N} u^2 \log u^2 dx = -\infty$.

The initial work to deal with this difficulty is due to Cazenave in [16], where the author considered the following logarithmic Schrödinger equation

$$iu_t + \Delta u + u \log u^2 = 0, \ (t, x) \in \mathbb{R} \times \mathbb{R}^N,$$

in the space $W \triangleq \{u \in H^1(\mathbb{R}^N) : \int_{\mathbb{R}^N} |u^2 \log u^2| dx < \infty\}$ with a suitable Luxemburg norm. Speaking it more clearly, by introducing the *N*-function

$$A(s) = \begin{cases} -s^{2} \log s^{2}, & 0 \le s \le e^{-3}, \\ 3s^{2} + 4e^{-3}s - e^{-6}, & s \ge e^{-3}, \end{cases}$$
(1.5)
$$\|u\|_{A} = \inf \left\{ \gamma > 0 : \int_{\mathbb{R}^{N}} A\left(\frac{|u|}{\gamma}\right) dx \le 1 \right\},$$

and $(\| \cdot \|_{H^1(\mathbb{R}^N)} + \| \cdot \|_A)$ as the Luxemburg norm, then the author obtained the existence of infinitely many critical points for the variational functional

$$\mathcal{J}(u) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \int_{\mathbb{R}^N} u^2 \log u^2 dx, \ u \in W,$$

on the set $\Sigma \triangleq \{u \in W : \int_{\mathbb{R}^N} u^2 dx = 1\}$. Very recently, Alves and his collaborators [3,4,8] have used the decomposition

$$\hat{F}_2(s) - \hat{F}_1(s) = \frac{1}{2}s^2 \log s^2, \ s \in \mathbb{R},$$
(1.6)

where,

$$\hat{F}_1(s) = \begin{cases} 0, & s \le 0, \\ -\frac{1}{2}s^2 \log s^2, & 0 < s < \delta, \\ -\frac{1}{2}s^2 \left[\log \delta^2 + 3 \right] + 2\delta s - \frac{1}{2}\delta^2, & s \ge \delta, \end{cases}$$

and

$$\hat{F}_2(s) = \begin{cases} 0, & |s| \le \delta, \\ \frac{1}{2}|s|^2 \log\left(|s|^2/\delta^2\right) + 2\delta|s| - \frac{3}{2}|s|^2 - \frac{1}{2}\delta^2, & |s| \ge \delta, \end{cases}$$

for some sufficiently small $\delta > 0$, and introduced the Orlicz space

$$L^{\hat{F}_1}(\mathbb{R}^N) = \left\{ u \in L^1_{\text{loc}}(\mathbb{R}^N) : \int_{\mathbb{R}^N} \hat{F}_1\left(\frac{|u|}{\gamma}\right) dx < +\infty \text{ for some } \gamma > 0 \right\},\$$

then studied some different types of logarithmic Schrödinger equation in the space $H^1(\mathbb{R}^N) \cap L^{\tilde{F}_1}(\mathbb{R}^N)$. Of course, there are various techniques in the literature, see e.g. [5–7,18,19,30,34,43, 46,50,55] and the references therein even if these ones are far to be exhaustive, to find some other interesting works on logarithmic Schrödinger equations.

On the other hand, one can deal with the case $\lambda \in \mathbb{R}$ is unknown. In this situation, $\lambda \in \mathbb{R}$ appears as a Lagrange multiplier and the L^2 -norm of solution is prescribed. From the physical point of view, the research holds particular significance as it accounts for the conservation of mass. Additionally, it provides valuable insights into the dynamic properties of standing waves in Eq. (1.4), such as stability or instability [15,17]. In [35], combining a minimax approach and compactness argument, Jenajean contemplated the existence of solutions for the following Schrödinger problem

$$\begin{cases} -\Delta u + \lambda u = g(u) \text{ in } \mathbb{R}^N, \\ \int\limits_{\mathbb{R}^N} |u|^2 dx = a^2 > 0. \end{cases}$$
(1.7)

Later on, there are some complements and generalizations in [37]. In [48], for $g(t) = \mu |t|^{q-2}t + |t|^{p-2}t$ with $2 < q \le 2 + \frac{4}{N} \le p < 2^*$, Soave considered the existence of solutions for problem (1.7), where $2^* = \frac{2N}{N-2}$ if $N \ge 3$ and $2^* = \infty$ if N = 2. For this type of combined nonlinearities, Soave [49] proved the existence of ground state and excited solutions when $p = 2^*$. For more interesting results for problem (1.7), we refer the reader to [13,36,38,40,56] and the references therein.

In reality, the *p*-Laplacian operator in Eq. (1.1) appears in many nonlinear problems, see [29,33,42] for example, in fluid dynamics, the shear stress $\vec{\tau}$ combined with the velocity gradient ∇u of the fluid corresponds to the manner that

$$\vec{\tau} = r |\nabla u|^{p-2} \nabla u.$$

The fluid is dilatant, pseudoplastic or Newtonian when p > 2, p < 2 or p = 2, respectively. Therefore, the equation governing the motion of the fluid includes the *p*-Laplacian. Moreover, such an operator also appears in the study of flows through porous media (when p = 3/2), non-linear elasticity (when $p \ge 2$) and glaciology (when $p \in (1, 4/3]$).

In light of the physical background of *p*-Laplacian operator, motivated by the prescribed L^2 norm solution for Eq. (1.4), it is interesting to investigate the *p*-Laplacian equation under the L^p -mass constraint and regard the frequency $\lambda \in \mathbb{R}$ as a Lagrange multiplier. Up to the best knowledge of us, there exist very few articles on this topic. Wang et al. [53] established the existence of solutions for the problem

$$\begin{cases} -\Delta_p u + |u|^{p-2}u = \lambda u + |u|^{s-1}u \text{ in } \mathbb{R}^N, \\ \int |u|^2 dx = a^2 > 0. \\ \mathbb{R}^N \end{cases}$$

when a > 0 is sufficiently small, where $\max\left\{1, \frac{2N}{N+2}\right\} and <math>s \in \left(\frac{N+2}{N}p, p^*\right)$ with $p^* = \frac{Np}{N-p}$. In [61], the authors studied the *p*-Laplacian equation with a L^p -norm constraint

$$\begin{cases} -\Delta_p u = \lambda |u|^{p-2} u + \mu |u|^{q-2} u + g(u) \text{ in } \mathbb{R}^N, \\ \int\limits_{\mathbb{R}^N} |u|^p \mathrm{d}x = a^p, \end{cases}$$

where $1 and <math>g \in C^0(\mathbb{R}, \mathbb{R})$ is odd and L^p -supercritical. For the suitable μ , they obtained several existence results including the existence of infinitely many solutions by Schwarz rearrangement technique, Ekeland variational principle and the Fountain theorem. There exist some other similar results for the *p*-Laplacian equation prescribed a L^p -norm, see e.g. [24,25,59].

Whereas, as far as we are concerned, there seems no related results for the *p*-Laplacian equation with a logarithmic nonlinearity under the L^p -mass constraint, and so one of the aims in the present article is to fulfill the blank. Motivated by [2,8,10], we shall derive the multiplicity and concentrating behavior of positive solutions for a logarithmic *p*-Laplacian equation (1.1) under the constraint (1.2).

Now, we can state the first main result as follows.

Theorem 1.1. Let $2 \le p < N$ and $(V_1) - (V_2)$. Then, there exists a $\varepsilon^* > 0$ such that (1.1)-(1.2) possesses at least l different couples of weak solutions $(u_{\varepsilon}^j, \lambda_{\varepsilon}^j) \in W^{1,p}(\mathbb{R}^N) \times \mathbb{R}$ for all $\varepsilon \in (0, \varepsilon^*)$ with $u_{\varepsilon}^j(x) > 0$ for every $x \in \mathbb{R}^N$ and $\lambda_{\varepsilon}^j < 0$, where $j \in \{1, 2, \dots, l\}$. Moreover, each u_{j}^{ε} has a maximum point $z_{\varepsilon}^j \in \mathbb{R}^N$ such that $V(z_{\varepsilon}^j) \to V(x^j) = V_0$ as $\varepsilon \to 0^+$. Besides, there exist two constants $C_0^j > 0$ and $c_0^j > 0$ satisfying

$$u_{\varepsilon}^{j} \leq C_{0}^{j} \exp\left(-c_{0}^{j} \frac{|x-z_{\varepsilon}^{j}|}{\varepsilon}\right)$$

for all $\varepsilon \in (0, \varepsilon^*)$ and $x \in \mathbb{R}^N$.

Remark 1.2. A recent work by Alves and Ji we would like to mention here is the paper [8], where the authors contemplated the following Schrödinger problem with logarithmic nonlinearity

$$\begin{cases} -\varepsilon^2 \Delta u + V(x)u = \lambda u + u \log |u|^2 \text{ in } \mathbb{R}^N, \\ \int_{\mathbb{R}^N} |u|^2 dx = a^2 \varepsilon^N. \end{cases}$$
(1.8)

Here the potential V satisfies the assumptions

$$(\hat{V}_1) \quad V \in \mathcal{C}^0(\mathbb{R}^N) \text{ and } -1 \le V_0 = \inf_{x \in \mathbb{R}^N} V(x) < V_\infty = \lim_{|x| \to +\infty} V(x) < +\infty; (\hat{V}_2) \quad M = \{x \in \mathbb{R}^N : V(x) = V_0\} \text{ and } M_{\hat{\delta}} = \{x \in \mathbb{R}^N : \operatorname{dist}(x, M) \le \hat{\delta}\} \text{ for some } \hat{\delta} > 0$$

They derived that, given a $\hat{\delta} > 0$, there exist $\hat{a} > 0$ and $\hat{\varepsilon} > 0$ such that problem (1.8) possesses at lease $\operatorname{cat}_{M_{\hat{\delta}}}(M)$ couples of solutions for all $a > \hat{a}$ and $\varepsilon \in (0, \hat{\varepsilon})$, where the concentration has also been studied but the property of exponential decay of solutions is absent. In contrast to it, there are three main contributions in the present paper which are exhibited below

(I) We just suppose that $V_0 > -\infty$ in (V_1) instead of $V_0 \ge -1$ in (\hat{V}_1) ;

(II) There is no restriction on the mass a > 0 in Theorem 1.1, while the results in [8] hold true provided that the mass a > 0 is sufficiently large;

(III) The *p*-Laplacian operator appearing in Eq. (1.1) is non-homogeneous which reveals that the calculations would be more complicated and technical. For instance, as a counterpart of (1.6), it is nontrivial to construct some functions F_1 and F_2 such that

$$F_2(s) - F_1(s) = \frac{1}{p} |s|^p \log |s|^p, \ \forall s \in \mathbb{R}.$$
(1.9)

Moreover, we shall make some efforts to investigate the property of exponential decay of obtained solutions which should be regarded as a replenishment. As a consequence, we could never repeat the approaches adopted in [8] to conclude the proof of Theorem 1.1.

We note that the considerations of Eq. (1.1) can date back to the studies of semiclassical problems for Schrödinger equations, the reader could refer to [12,14] for some detailed survey on such topic which comes from the pioneering research work by Floer and Weinstein [32]. Soon afterwards, this topic on different types of Schrödinger equations has been investigated

extensively under several distinct hypotheses on the potential and the nonlinearity, see e.g. [11, 22,23,27,28,39,44,54] and the references therein. Hence, it permits us to follow the effective procedures in the literature to handle the *p*-Laplacian problems with $2 \le p < N$ and logarithmic nonlinearity in this paper.

Performing the scaling $v(x) = u(\varepsilon x)$, one could observe that, to consider (1.1)-(1.2), it is equivalent to study the problem

$$\begin{cases} -\Delta_p v + V(\varepsilon x)|v|^{p-2}v = \lambda|v|^{p-2}v + |v|^{p-2}v\log|v|^p \text{ in } \mathbb{R}^N,\\ \int_{\mathbb{R}^N} |v|^p dx = a^p. \end{cases}$$
(1.10)

In other words, if the couple (v, λ) is a (weak) solution of Problem (1.10), then (u, λ) is a solution of (1.1)-(1.2), where $v(x) = u(\varepsilon x)$ for all $x \in \mathbb{R}^N$. Let $v = \sigma w$ with some $\sigma > 0$, we shall observe that the couple (v, λ) is a weak solution to Problem (1.10) if and only if (w, λ) is a weak solution to problem below

$$\begin{cases} -\Delta_p w + [V(\varepsilon x) - \log \sigma^p] |w|^{p-2} w = \lambda |w|^{p-2} w + |w|^{p-2} w \log |w|^p \text{ in } \mathbb{R}^N, \\ \int_{\mathbb{R}^N} |w|^p dx = a^p \sigma^{-p}. \end{cases}$$
(1.11)

At this stage, we deduce that if one wants to study problems (1.1)-(1.2), it would be enough to deal with Problem (1.11). Since $\sigma > 0$ is arbitrary, we are able to find a sufficiently small $\sigma > 0$ such that

$$V(\varepsilon x) - \log \sigma^p \ge -1, \ \forall x \in \mathbb{R}^N,$$

since $V_0 = \inf_{x \in \mathbb{R}^N} V(x) > -\infty$ by (V_1) and $a^p \sigma^{-p} > \tilde{a}$, where $\tilde{a} > 0$ is a fixed constant.

Owing to the above discussions, to reach Theorem 1.1, we just need to prove the following result.

Theorem 1.3. Let $2 \le p < N$ and $(\hat{V}_1) - (V_2)$. Then, there exist $a^* > 0$ and $\varepsilon^* > 0$ such that (1.1)-(1.2) has at least l different couples of weak solutions $(u_{\varepsilon}^j, \lambda_{\varepsilon}^j) \in W^{1,p}(\mathbb{R}^N) \times \mathbb{R}$ for all $a > a^*$ and $\varepsilon \in (0, \varepsilon^*)$ with $u_{\varepsilon}^j(x) > 0$ for every $x \in \mathbb{R}^N$ and $\lambda_{\varepsilon}^j < 0$, where $j \in \{1, 2, \dots, l\}$. Moreover, each u_{ε}^j admits a maximum point $z_{\varepsilon}^j \in \mathbb{R}^N$ such that $V(z_{\varepsilon}^j) \to V(x^j) = V_0$ as $\varepsilon \to 0^+$. Besides, there exist two constants $C_0^j > 0$ and $c_0^j > 0$ satisfying

$$u_{\varepsilon}^{j} \leq C_{0}^{j} \exp\left(-c_{0}^{j} \frac{|x-z_{\varepsilon}^{j}|}{\varepsilon}\right)$$

for all $\varepsilon \in (0, \varepsilon^*)$ and $x \in \mathbb{R}^N$.

Now, we shall turn to investigate the existence of normalized solutions for logarithmic *p*-Laplacian equations with mass subcritical and supercritical nonlinearities. First of all, we recall the Gagliardo-Nirenberg inequality (see e.g. [1,57]), for every $p < s < p^*$, there exists an optimal constant $\mathbb{C}_{N,p,s} > 0$ depending only on *N*, *p* and *s* such that

$$\|u\|_{L^{s}(\mathbb{R}^{N})} \leq \mathbb{C}_{N,p,s} \|\nabla u\|_{L^{p}(\mathbb{R}^{N})}^{\beta_{s}} \|u\|_{L^{p}(\mathbb{R}^{N})}^{1-\beta_{s}}, \ \forall u \in W^{1,p}(\mathbb{R}^{N}),$$
(1.12)

where

$$\beta_s \triangleq N\left(\frac{1}{p} - \frac{1}{s}\right) = \frac{N(s-p)}{ps}.$$
(1.13)

Due to (1.12), one observes that

$$\bar{p} = p + \frac{p^2}{N}$$

is the L^p -critical exponent with respect to p-Laplacian equation. Indeed, we focus on establishing the existence and multiplicity of positive solutions for the following p-Laplacian type problems

$$\begin{cases} -\Delta_p u = \lambda |u|^{p-2} u + |u|^{p-2} u \log |u|^p + \mu |u|^{q-2} u \text{ in } \mathbb{R}^N, \\ \int\limits_{\mathbb{R}^N} |u|^p dx = a^p, \end{cases}$$
(1.14)

where $a > 0, \mu > 0$ and $q \in (p, \bar{p}) \cup (\bar{p}, p^*)$.

In order to contemplate the Problem (1.14) involving a class of pure-power type mass subcritical and supercritical nonlinearities, we shall continue to establish the existence of global minimizer of a minimization problem, namely

$$m(a) = \inf_{u \in S(a)} J(u),$$

where the variational functional $J: X \to \mathbb{R}$ is given by

$$J(u) = \frac{1}{p} \int_{\mathbb{R}^{N}} \left(|\nabla u|^{p} + |u|^{p} \right) dx - \frac{1}{p} \int_{\mathbb{R}^{N}} |u|^{p} \log |u|^{p} dx - \frac{\mu}{q} \int_{\mathbb{R}^{N}} |u|^{q} dx$$
(1.15)

and the constrained set

$$S(a) = \left\{ u \in X : \int_{\mathbb{R}^N} |u|^p dx = a^p \right\}.$$

Here a > 0 and the space X can be found in Section 2 below.

The main results for the Problem (1.14) involving mass-subcritical nonlinearities can be stated as follows.

Theorem 1.4. Suppose that $2 \le p < N$, $\mu > 0$ and $p < q < \overline{p}$. Then, there is an $a_* > 0$ such that, for all fixed $a > a_*$, m(a) admits a minimizer $u \in S(a)$ which is positive and radially symmetric and decreasing in r = |x|. Moreover, there is $a \lambda \in \mathbb{R}$ such that $(u, \lambda) \in X \times \mathbb{R}$ solves Problem (1.14).

Remark 1.5. It should be stressed here that Theorem 1.4 partially generalizes [47, Theorem 1.1]. Nevertheless, one could never simply repeat the arguments explored in it to arrive at our result since it needs some efforts to construct a suitable *N*-function like (1.5) in the *p*-Laplacian setting. In order to avoid this obstacle, we continue to make use of the decomposition in (1.9) which should be regarded as one of novelties in this article. Whereas, the biggest challenge in the proof of Theorem 1.4 is the lack of compactness. Explaining it more clearly, the imbedding $W_r \hookrightarrow L^2(\mathbb{R}^N)$ in [47] is compact, see e.g. [16, Proposition 3.1], but we indeed can not conclude the compact imbedding $X_r \hookrightarrow L^p(\mathbb{R}^N)$ in advance. To overcome this difficulty, we shall introduce some new analytic tricks to recover the desired compactness, see Lemma 4.4 for instance. Moreover, the reader would observe that the case $\mu = 0$ in Theorem 1.4 acts as a special one of Theorem 3.1 below. Alternatively, we successfully put forward a totally different argument to look for a minimizer of m(a).

Finally, we begin paying attention to the case $p + \frac{p^2}{N} < q < p^*$ which leads to the Problem (1.14) a mass-supercritical one and prove the following result.

Theorem 1.6. Suppose that $2 \le p < N$ and $p + \frac{p^2}{N} < q < p^*$, there exist $\underline{a}^* > 0$ and $\underline{\mu}^* > 0$ such that, for all $a > \underline{a}^*$ and $\mu \in (0, \underline{\mu}^*)$, Problem (1.14) has a couple of weak solution $(\underline{u}^*, \underline{\lambda}^*) \in X \times \mathbb{R}$ with $\underline{u}^*(x) > 0$ for all $x \in \mathbb{R}^N$.

Given a $u \in S(a)$, then $u_t(\cdot) \triangleq t^{\frac{N}{p}}u(t\cdot)$ for all t > 0 and hence, it is elementary to arrive at the calculations

$$J(u_t) = \frac{t^p}{p} \int_{\mathbb{R}^N} |\nabla u|^p dx + \frac{a^p (1 - N \log t)}{p} - \frac{1}{p} \int_{\mathbb{R}^N} |u|^p \log |u|^p dx - \frac{\mu}{q} t^N \left(\frac{q}{p} - 1\right) \int_{\mathbb{R}^N} |u|^q dx$$
$$\to -\infty$$

as $t \to +\infty$ which indicates that $m(a) = -\infty$ for all a > 0. To get around this obstacle, in a similar spirit of [13], one may depend on the following Pohožaev manifold

$$\mathcal{P}(a) = \{ u \in S(a) : P(u) = 0 \},\$$

where the variational functional $P: S(a) \to \mathbb{R}$ is defined by

$$P(u) = \int_{\mathbb{R}^N} |\nabla u|^p dx - \frac{N}{p} a^p - \mu N\left(\frac{1}{p} - \frac{1}{q}\right) \int_{\mathbb{R}^N} |u|^q dx.$$

According to the discussions in Section 2 below, we can conclude that \mathcal{P} is a natural constraint since $P(u) \equiv 0$ provided that $u \in X$ is a nontrivial weak solution of the first equation in Problem (1.14). Unfortunately, we fail to argue as [47] to consider the following minimization problem

$$m_p(a) = \inf_{u \in \mathcal{P}(a)} J(u)$$

since it seems impossible to get the compactness of its corresponding minimizing sequence. Actually, we even cannot make sure that $m_p(a) \le 0$. Thereby, it differs evidently from the counterparts in [47]. Besides, we also try to study the following problem

$$m_p^{\mathcal{R}}(a) = \inf_{u \in S(a) \cap \{ \|\nabla u\|_{L^p(\mathbb{R}^N)} < \mathcal{R} \}} J(u)$$

combined with the Ekeland variational principle. Although one can deduce that $m_p^{\mathcal{R}}(a) < 0$ for some suitable $\mathcal{R} > 0$, the absence of the monotone property with respect to $m_p^{\mathcal{R}}(a)$ results in the lack of compactness and it is still hard to show that $m_p^{\mathcal{R}}(a)$ can be attained.

In consideration of the explanations exhibited above, motivated by the ideas introduced in [9], we shall rely heavily on the so-called truncation argument. Let us introduce it step by step. For every $\mathcal{R} > 0$ and $p < \bar{q} < p + \frac{p^2}{N}$, we define the auxiliary function $f_{\mathcal{R}} : \mathbb{R} \to \mathbb{R}$ given by

$$f_{\mathcal{R}}(t) = \begin{cases} |t|^{q-2}t, & |t| \leq \mathcal{R}, \\ \mathcal{R}^{q-\bar{q}}|t|^{\bar{q}-2}t, & |t| \geq \mathcal{R}. \end{cases}$$

Using the function $f_{\mathcal{R}}$, we then contemplate the following auxiliary problem

$$\begin{cases} -\Delta_{p}u = \lambda |u|^{p-2}u + |u|^{p-2}u \log |u|^{p} + \mu f_{\mathcal{R}}(u) \text{ in } \mathbb{R}^{N}, \\ \int_{\mathbb{R}^{N}} |u|^{p} dx = a^{p}, \\ \mathbb{R}^{N} \end{cases}$$
(1.16)

whose energy functional $J_{\mathcal{R}}: X \to \mathbb{R}$ is given by

$$J_{\mathcal{R}}(u) = \frac{1}{p} \int_{\mathbb{R}^N} \left(|\nabla u|^p + |u|^p \right) dx - \frac{1}{p} \int_{\mathbb{R}^N} |u|^p \log |u|^p dx - \mu \int_{\mathbb{R}^N} F_{\mathcal{R}}(u) dx,$$

where and in the sequel $F_{\mathcal{R}}(t) = \int_0^t f_{\mathcal{R}}(s) ds$. Due to the definition of $f_{\mathcal{R}}$, one has that

$$|f_{\mathcal{R}}(t)| \le \mathcal{R}^{q-\bar{q}}|t|^{\bar{q}-1}, \ \forall t \in \mathbb{R}.$$
(1.17)

When the constant $\mathcal{R} > 0$ is fixed, it follows from (1.17) that $f_{\mathcal{R}}$ has a L^p -subcritical growth because $p < \bar{q} < p + \frac{p^2}{N}$. In other words, adopting Theorem 1.4, we immediately have the results below.

Corollary 1.7. Suppose that $2 \le p < N$, $\mu > 0$ and $p < \bar{q} < p + \frac{p^2}{N}$. Then, for every fixed $\mathcal{R} > 0$, there is an $a^* > 0$ independent of \mathcal{R} and μ such that, for all fixed $a > a^*$, Problem (1.16) admits a couple of weak solution $(u_{\mathcal{R}}^*, \lambda_{\mathcal{R}}^*) \in X \times \mathbb{R}$ with $u_{\mathcal{R}}^*(x) > 0$ for all $x \in \mathbb{R}^N$.

With Corollary 1.7 in hands, the reader is invited to see that if $u_{\mathcal{R}}^* \in X$ is a solution of Problem (1.16) with $|u_{\mathcal{R}}^*|_{\infty} \leq \mathcal{R}$, then $u_{\mathcal{R}}^*$ is a solution for Problem (1.14). Have this in mind, our main goal is to deduce that given an $\mathcal{R} > 0$, there are $\underline{a}^* > 0$ (independent of \mathcal{R} and μ) and $\underline{\mu}^* = \mu^*(\mathcal{R}) > 0$ such that if $a > \underline{a}^*$ and $\mu \in (0, \mu^*)$, then $|u_{\mathcal{R}}^*|_{\infty} \leq \mathcal{R}$.

We remark that, d'Avenia, Montefusco and Squassina [18] handled the existence of infinitely many solutions for a class of logarithmic Schrödinger equations. The authors pointed out their multiplicity results are also adapted, using [21], to the following logarithmic *p*-Laplacian equation

$$-\Delta_p u = \lambda |u|^{p-2} u + |u|^{p-2} u \log |u|^p, \ u \in W^{1,p}(\mathbb{R}^N),$$
(1.18)

where $\lambda \in \mathbb{R}$ is a fixed constant. On the other hand here we mainly focus on existence of families of solutions concentrating around local minima of *V* in the semiclassical limit $\varepsilon \to 0$.

Again the results in Theorems 1.3, 1.4 and 1.6 are new under the *p*-Laplacian settings with the logarithmic nonlinearity. The striking novelty is the correct setting of functional space in which we can treat the problems variationally. Unfortunately, we cannot deal with the case 1 so far ant it remains open, see Lemma 2.2 below. In addition, there are some other technical calculations due to the*p*-Laplacian operator in the proofs of the main results.

The outline of the paper is organized as follows. In Section 2, we mainly exhibit some preliminary results. Sections 3 and 4 are devoted to the non-autonomous and autonomous logarithmic p-Laplacian equations, respectively. Finally, there are some further comments in Section 5.

Notations. From now on in this paper, otherwise mentioned, we use the following notations:

- C, C_1, C_2, \cdots denote any positive constant, whose value is not relevant.
- Let $(Z, \|\cdot\|_Z)$ be a Banach space with its dual space $(Z^*, \|\cdot\|_{Z^*})$.
- $|\cdot|_p$ denotes the usual norm of the Lebesgue measurable space in \mathbb{R}^N , for all $p \in [1, +\infty]$.
- $o_n(1)$ denotes the real sequence with $o_n(1) \to 0$ as $n \to +\infty$.
- " \rightarrow " and " \rightarrow " stand for the strong and weak convergence in the related function spaces, respectively.

2. Variational setting and preliminaries

In this section, we would like to recommend some preliminary results. First of all, let us introduce some fundamental concepts and properties concerning the Orlicz spaces. For the more details, please refer to [45] for example.

Definition 2.1. An *N*-function is a continuous function $\Phi : \mathbb{R} \to [0, +\infty)$ that satisfies the following conditions:

(i) Φ is a convex and even function;

(ii)
$$\Phi(t) = 0 \iff t = 0;$$

(iii) $\lim_{t \to 0} \frac{\Phi(t)}{t} = 0$ and $\lim_{t \to \infty} \frac{\Phi(t)}{t} = +\infty$.

We say that an N-function Φ satisfies the Δ_2 -condition, denoted by $\Phi \in (\Delta_2)$, if

$$\Phi(2t) \le k\Phi(t), \ \forall t \ge t_0,$$

for some constants k > 0 and $t_0 \ge 0$.

The conjugate function Φ associated with Φ is obtained through the Legendre's transformation, defined as

$$\tilde{\Phi}(s) = \max_{t \ge 0} \{st - \Phi(t)\}, \text{ for } s \ge 0.$$

It can be shown that $\tilde{\Phi}$ is also an *N*-function. The functions Φ and $\tilde{\Phi}$ are mutually complementary that is, $\tilde{\tilde{\Phi}} = \Phi$.

For an open set $\Omega \subset \mathbb{R}^N$, we define the Orlicz space associated with the *N*-function Φ as follows

$$L^{\Phi}(\Omega) = \left\{ u \in L^{1}_{\text{loc}}(\Omega) : \int_{\Omega} \Phi\left(\frac{|u|}{\gamma}\right) dx < +\infty, \text{ for some } \gamma > 0 \right\},\$$

which is a Banach space endowed with the Luxemburg norm given by

$$||u||_{\Phi} = \inf \left\{ \gamma > 0 : \int_{\Omega} \Phi\left(\frac{|u|}{\gamma}\right) dx \le 1 \right\}.$$

Associated with the Orlicz Spaces, there also holds the Hölder and Young type inequalities, namely

$$st \le \Phi(t) + \tilde{\Phi}(s), \quad \forall s, t \ge 0$$

and

$$\left| \int_{\Omega} uv dx \right| \le 2 \|u\|_{\Phi} \|v\|_{\tilde{\Phi}}, \text{ for } \forall u \in L^{\Phi}(\Omega) \text{ and } \forall v \in L^{\bar{\Phi}}(\Omega).$$

The space $L^{\Phi}(\Omega)$ is reflexive and separable provided that $\Phi, \tilde{\Phi} \in (\Delta_2)$. Moreover, the Δ_2 condition implies that

$$L^{\Phi}(\Omega) = \left\{ u \in L^{1}_{\text{loc}}(\Omega) : \int_{\Omega} \Phi(|u|) dx < +\infty \right\}$$

and

$$u_n \to u \text{ in } L^{\Phi}(\Omega) \iff \int_{\Omega} \Phi(|u_n - u|) \, dx \to 0.$$

We then recall an significant relation involving *N*-functions that will be adopted later. Let Φ be an *N*-function of C^1 class and $\tilde{\Phi}$ is its conjugate function. Suppose that

$$1 < l \le \frac{\Phi'(t)t}{\Phi(t)} \le m, \ t \ne 0, \tag{2.1}$$

then Φ , $\tilde{\Phi} \in (\Delta_2)$. Finally, we consider the functions

Journal of Differential Equations 421 (2025) 1-49

L. Shen and M. Squassina

$$\xi_0(t) = \min\{t^l, t^m\}$$
 and $\xi_1(t) = \max\{t^l, t^m\}, t \ge 0$,

it is possible to verify that, using (2.1), the function Φ satisfies the inequality below

$$\xi_0\left(\|u\|_{\Phi}\right) \le \int_{\mathbb{R}^N} \Phi(u) \le \xi_1\left(\|u\|_{\Phi}\right), \ \forall u \in L^{\Phi}(\Omega).$$

$$(2.2)$$

Inspired by [3,4,8], we define the functions F_1 and F_2 as follows

$$F_{1}(s) = \begin{cases} F_{1}(-s), & s \leq 0, \\ -\frac{1}{p}s^{p}\log s^{p}, & 0 < s < (p-1)\delta, \\ -\frac{1}{p}s^{p} \Big[\log\left((p-1)\delta\right)^{p} + p + 1\Big] + p\delta s^{p-1} - \frac{1}{p(p-1)} \big((p-1)\delta\big)^{p}, & s \geq (p-1)\delta, \end{cases}$$

and

$$F_2(s) = \begin{cases} 0, & |s| \le (p-1)\delta, \\ \frac{1}{p} |s|^p \log\left(|s|^p / ((p-1)\delta)^p\right) + p\delta|s|^{p-1} - \frac{p+1}{p} |s|^p - \frac{1}{p(p-1)} \left((p-1)\delta\right)^p, \\ & |s| \ge (p-1)\delta, \end{cases}$$

where $\delta > 0$ is sufficiently small but fixed, then we reach the decomposition (1.9). Moreover, F_1 and F_2 satisfy the following properties:

- (**P**₁) F_1 is even with $F'_1(s)s \ge 0$ and $F_1(s) \ge 0$ for all $s \in \mathbb{R}$. Furthermore, $F_1 \in \mathcal{C}^1(\mathbb{R}, \mathbb{R})$ is convex if $\delta \approx 0^+$;
- (**P**₂) $F_2 \in \mathcal{C}^1(\mathbb{R}, \mathbb{R}) \cap \mathcal{C}^2((\delta, +\infty), \mathbb{R})$ and for each $\tilde{q} \in (p, p^*)$, there exists a $C_{\tilde{q}} > 0$ such that

$$|F'_2(s)| \le C_{\tilde{q}}|s|^{\tilde{q}-1}, \quad \forall s \in \mathbb{R};$$

(**P**₃) $s \mapsto \frac{F'_2(s)}{s^{p-1}}$ is a nondecreasing function for s > 0 and a strictly increasing function for $s > \delta$; (**P**₄) $\lim_{s \to \infty} \frac{F'_2(s)}{s^{p-1}} = \infty$.

As a counterpart of the results explored in [3,4,8], we demonstrate the following result which is nontrivial in contrast to the cited papers.

Lemma 2.2. The function F_1 is an N-function. Moreover, if $2 \le p < N$, it holds that $F_1, \tilde{F}_1 \in (\Delta_2)$.

Proof. Exploiting some elementary calculations, one could easily certify that F_1 satisfies (I)-(III) of Definition 2.1. To arrive at the proof, we shall verify that F_1 satisfies a similar relation in (2.1) and so it reveals that F_1 , $\tilde{F}_1 \in (\Delta_2)$. Firstly, it is easy to see that

$$F'_1(s) = \begin{cases} -(1 + \log s^p)s^{p-1}, & 0 < s < (p-1)\delta, \\ -s^{p-1} \left[\log \left((p-1)\delta \right)^p + p + 1 \right] + p(p-1)\delta s^{p-2}, & s \ge (p-1)\delta. \end{cases}$$

Next, we shall analyze the cases $0 < s < (p-1)\delta$ and $s \ge (p-1)\delta$ separately.

Case 1. $0 < s < (p-1)\delta$.

In this case, it is simple to calculate that

$$\frac{F_1'(s)s}{F_1(s)} = p + \frac{1}{\log s},$$

which indicates that there is an $l_1 > 1$ such that

$$1 < l_1 \le \frac{F_1'(s)s}{F_1(s)} \le m_1 \triangleq \sup_{0 < s < \delta} \left(p + \frac{1}{\log s} \right) \le p,$$

for some sufficiently small $\delta > 0$.

Case 2. $s \ge (p-1)\delta$.

In this case, we continue to calculate that

$$\frac{F_1'(s)s}{F_1(s)} = \frac{-s^p \left[\log\left((p-1)\delta\right)^p + p + 1\right] + p(p-1)\delta s^{p-1}}{-\frac{1}{p}s^p \left[\log\left((p-1)\delta\right)^p + p + 1\right] + p\delta s^{p-1} - \frac{1}{p(p-1)}\left((p-1)\delta\right)^p}$$

From which, we derive that $\sup_{s \ge (p-1)\delta} \frac{F'_1(s)s}{F_1(s)} \le p$ since for all $s \ge (p-1)\delta$, there holds

$$\frac{F_1'(s)s}{F_1(s)} \le \frac{-s^p \Big[\log \big((p-1)\delta \big)^p + p + 1 \big] + p(p-1)\delta s^{p-1} + \Big[p\delta s^{p-1} - \frac{1}{p-1} \big((p-1)\delta \big)^p \Big]}{-\frac{1}{p} s^p \Big[\log \big((p-1)\delta \big)^p + p + 1 \Big] + p\delta s^{p-1} - \frac{1}{p(p-1)} \big((p-1)\delta \big)^p \Big]}$$

Obviously, one can deduce that

$$\lim_{s \to +\infty} \frac{F_1'(s)s}{F_1(s)} = p \text{ and } \frac{F_1'(s)s}{F_1(s)} > p - 1, \ \forall s > 0,$$

and so we obtain

$$p-1 < \inf_{s>0} \frac{F_1'(s)s}{F_1(s)}.$$

The last inequality together with p > 2 guarantees the existence of an $l \in (1, 2)$ such that

$$1 < l \le \frac{F_1'(s)s}{F_1(s)} \le p, \ \forall s > 0.$$

Since F_1 is an even function, then the inequality holds true for any $s \neq 0$. The proof is completed. \Box

Replacing Φ and Ω in the above discussions with F_1 and \mathbb{R}^N , respectively, we conclude the Orlicz Space $L^{F_1}(\mathbb{R}^N)$ and it is standard to prove the following result.

Corollary 2.3. The functional $\Theta: L^{F_1}(\mathbb{R}^N) \to \mathbb{R}$ given by $u \mapsto \int_{\mathbb{R}^N} F_1(u) dx$ is of class $\mathcal{C}^1(L^{F_1}(\mathbb{R}^N))$ with

$$\Theta'(u)v = \int_{\mathbb{R}^N} F'_1(u)v dx, \ \forall u, v \in L^{F_1}(\mathbb{R}^N),$$

where $L^{F_1}(\mathbb{R}^N)$ denotes the Orlicz space associated with F_1 endowed with the Laremburg norm $\|\cdot\|_{F_1}$.

In the sequel, in order to avoid the points $u \in W^{1,p}(\mathbb{R}^N)$ that satisfy $F_1(u) \notin L^1(\mathbb{R}^N)$, we should consider the work space $X = W^{1,p}(\mathbb{R}^N) \cap L^{F_1}(\mathbb{R}^N)$ throughout the paper equipped with the norm

$$\|\cdot\| \triangleq \|\cdot\|_{W^{1,p}(\mathbb{R}^N)} + \|\cdot\|_{F_1},$$

where $\|\cdot\|_{W^{1,p}(\mathbb{R}^N)}$ denotes the usual norm in $W^{1,p}(\mathbb{R}^N)$. Moreover, we denote the radially symmetric subsequence of X by X_r , namely $X_r = \{u \in X : u(x) = u(|x|)\}$ with the norm $\|\cdot\|$.

With the space X and (1.9) in hands, we can obtain the following Brézis-Lieb type lemma in the logarithmic setting.

Lemma 2.4. Let $\{u_n\}$ be a bounded sequence in X such that $u_n \to u$ a.e. in \mathbb{R}^N and $\{|u_n|^p \log |u_n|^p\}$ is a bounded sequence in $L^1(\mathbb{R}^N)$. Then, up to a subsequence if necessary,

$$\lim_{n\to\infty}\int_{\mathbb{R}^N} \left(|u_n|^p \log |u_n|^p - |u_n - u|^p \log |u_n - u|^p\right) dx = \int_{\mathbb{R}^N} |u|^p \log |u|^p dx.$$

Proof. Recalling (1.9), one has that

$$F_2(u_n) - F_1(u_n) = \frac{1}{p} |u_n|^p \log |u_n|^p.$$

Since $\{u_n\}$ is a bounded sequence in X, by property-(\mathbf{P}_2), it follows from [58, Lemma 1.32] that

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} [F_2(u_n) - F_2(u_n - u)] dx = \int_{\mathbb{R}^N} F_2(u) dx.$$

Similarly, we easily conclude that

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} [F_1(u_n) - F_1(u_n - u)] dx = \int_{\mathbb{R}^N} F_1(u) dx.$$

So, we can finish the proof of the lemma. \Box

Next, we shall introduce the Pohožaev identity for a class of logarithmic *p*-Laplacian equations in \mathbb{R}^N as follows.

Theorem 2.5. Let $2 \le p < N$. Suppose $u \in X$ to be a nontrivial weak solution of

$$-\Delta_p u = \lambda |u|^{p-2} u + |u|^{p-2} u \log |u|^p + \mu |u|^{q-2} u \text{ in } \mathbb{R}^N,$$
(2.3)

where $\lambda, \mu \in \mathbb{R}$ are constants and $p < q \leq p^*$. Then

$$\int_{\mathbb{R}^N} |\nabla u|^p dx = \frac{N}{p} \int_{\mathbb{R}^N} |u|^p dx + \mu N \left(\frac{1}{p} - \frac{1}{q}\right) \int_{\mathbb{R}^N} |u|^q dx.$$
(2.4)

Moreover, if in addition $u(x) \ge 0$ *for all* $x \in \mathbb{R}^N$ *, then* u(x) > 0 *for all* $x \in \mathbb{R}^N$ *.*

Proof. The proof is divided into the following three steps.

Step 1. $u \in L^{\infty}(\mathbb{R}^N) \cap \mathcal{C}^{1,\tau}_{\text{loc}}(\mathbb{R}^N)$ for some $\tau \in (0, 1)$.

We start by assuming that $u \ge 0$. For all L > 1, define $u_L = \min\{u, L\}$. Taking $\psi = u_L^{kp+1} \in X$ with $k \ge 0$ as a test function in (2.3), we obtain

$$\int_{\mathbb{R}^N} |\nabla u|^{p-2} \nabla u \nabla (u_L^{kN+1}) dx = \int_{\mathbb{R}^N} [F_2'(u) - F_1'(u) + (\lambda - 1)|u|^{p-2} u + \mu |u|^{q-2} u] u_L^{kp+1} dx.$$
(2.5)

It is easy to observe that

$$\begin{cases} \int \limits_{\mathbb{R}^{N}} |\nabla u|^{p-2} \nabla u \nabla (u_{L}^{kp+1}) dx = \frac{kp+1}{(k+1)^{p}} \int \limits_{\mathbb{R}^{N}} |\nabla (u_{L})^{k+1}|^{p} dx, \\ \int \limits_{\mathbb{R}^{N}} |u|^{p-2} u u_{L}^{kp+1} dx \ge \int \limits_{\mathbb{R}^{N}} |(u_{L})^{k+1}|^{p} dx. \end{cases}$$
(2.6)

Adopting property- (\mathbf{P}_1) and property- (\mathbf{P}_2) with $\tilde{q} = q$, there holds

$$F_{2}'(u)u - F_{1}'(u)u + \lambda|u|^{p} + \mu|u|^{q} \le (C_{q} + |\lambda| + |\mu|)|u|^{q} \triangleq C_{q,\lambda,\mu}|u|^{q} \text{ if } |u| \ge 1.$$
(2.7)

Without loss of generality, we shall suppose that $|u| \ge 1$. Combining (2.5), (2.6) and (2.7), we have

$$\left(\int_{\mathbb{R}^{N}} |u_{L}|^{(k+1)p^{*}}\right)^{\frac{p}{p^{*}}} \leq C_{p^{*}} \|(u_{L})^{k+1}\|_{W^{1,p}(\mathbb{R}^{N})}^{p} \leq C_{p^{*}}C_{q,\lambda,\mu}(k+1)^{p} \int_{\mathbb{R}^{N}} |u|^{q} u^{kp} dx$$
$$\leq C_{p^{*}}C_{q,\lambda,\mu}(k+1)^{p} \left(\int_{\mathbb{R}^{N}} |u|^{q} dx\right)^{\frac{q-p}{q}} \left(\int_{\mathbb{R}^{N}} |u|^{(k+1)q} dx\right)^{\frac{p}{q}} (2.8)$$

Letting $L \to +\infty$ in (2.8), we arrive at

$$\left(\int\limits_{\mathbb{R}^N} |u_L|^{(k+1)p^*}\right)^{\frac{p}{p^*}} \leq C_{p^*} C_{q,\lambda,\mu} (k+1)^p \mathbb{T}(u) \left(\int\limits_{\mathbb{R}^N} |u|^{(k+1)q} dx\right)^{\frac{p}{q}}$$

which is equivalent to

$$\left(\int_{\mathbb{R}^{N}} |u|^{(k+1)p^{*}}\right)^{\frac{1}{(k+1)p^{*}}} \leq C_{*}^{\frac{1}{k+1}}(k+1)^{\frac{1}{k+1}} \left(\int_{\mathbb{R}^{N}} |u|^{(k+1)q} dx\right)^{\frac{1}{(k+1)q}},$$
(2.9)

where the constant $C_* = C_{p^*}^p C_{q,\lambda,\mu}^p [\mathbb{T}(u)]^p > 0$ is independent of k. Let k = 0 in (2.9), it becomes

$$\left(\int_{\mathbb{R}^N} |u|^{q\varpi}\right)^{\frac{1}{q\varpi}} \leq C_* \left(\int_{\mathbb{R}^N} |u|^q dx\right)^{\frac{1}{q}},$$

where $\varpi = \frac{p^*}{q} \ge 1$. For $k + 1 = \varpi^m$ with $m \in \mathbb{N}^+$ in (2.9), it holds that

$$\left(\int_{\mathbb{R}^N} |u|^{\varpi^{m+1}\sigma} dx\right)^{\frac{1}{\varpi^{m+1}\sigma}} \le C_*^{\frac{1}{\varpi^m}} \varpi^{\frac{m}{\varpi^m}} \left(\int_{\mathbb{R}^N} |u|^{\varpi^m q} dx\right)^{\frac{1}{\varpi^m q}}$$

From it, proceeding this iteration procedure m times and multiplying these m + 1 formulas,

$$\left(\int\limits_{\mathbb{R}^N} |u|^{\varpi^{m+1}\sigma} dx\right)^{\frac{1}{\varpi^{m+1}\sigma}} \leq C_*^{\sum_{j=0}^m \frac{1}{\varpi^j}} \varpi^{\sum_{j=1}^m \frac{j}{\varpi^j}} \left(\int\limits_{\mathbb{R}^N} |u|^q dx\right)^{\frac{1}{q}}.$$

Since $\sum_{j=0}^{\infty} \frac{1}{\varpi^j} = \frac{\varpi}{\varpi-1}$ and $\sum_{j=1}^{\infty} \frac{j}{\varpi^j} = \frac{\varpi}{(\varpi-1)^2}$, then we could take the limit as $m \to +\infty$ to conclude that $u \in L^{\infty}(\mathbb{R}^N)$. When *u* changes sign, then it is enough to argue as before by contemplating once the positive part $u^+ \triangleq \max\{u, 0\}$ and once the negative part $u^- \triangleq \max\{-u, 0\}$ in place of *u* in the definition of u_L . As a result, we shall finish the verification of $u \in L^{\infty}(\mathbb{R}^N)$ for all nontrivial solution *u*. In addition, we could follow [26] to conclude that $u \in C_{\text{loc}}^{1,\tau}(\mathbb{R}^N)$ for some $\tau \in (0, 1)$.

Step 2. The nontrivial solution $u \in X$ satisfies (2.4).

We recall [20, Theorem 2] which is stated by Lemma A.1 in the Appendix and take $\mathcal{L}(x, s, \xi) = \frac{1}{p} |\xi|^p$ which is strictly convex in the variable $\xi \in \mathbb{R}^N$. Let $\varphi \in C_c^1(\mathbb{R}^N)$ be such that $0 \le \varphi \le 1$, $\varphi(x) = 1$ for all $|x| \le 1$, and $\varphi(x) = 0$ for all $|x| \ge 2$. Define

$$h(x) = \varphi\left(\frac{x}{k}\right) x \in \mathcal{C}^1(\mathbb{R}^N, \mathbb{R}^N), \text{ for all } k \in \mathbb{N}^+.$$

Note that if $h_j(x) = \varphi(\frac{x}{k})x_j$ for $j = 1, 2, \dots$, then

Journal of Differential Equations 421 (2025) 1-49

$$\begin{cases} D_i h_j(x) = D_i \varphi\left(\frac{x}{k}\right) \frac{x_j}{k} + \varphi\left(\frac{x}{k}\right) \delta_{ij}, \text{ for all } x \in \mathbb{R}^N, \ j = 1, 2, \cdots, \\ \operatorname{div} h(x) = D\varphi\left(\frac{x}{k}\right) \frac{x}{k} + N\varphi\left(\frac{x}{k}\right), \text{ for all } x \in \mathbb{R}^N, \end{cases}$$

where δ_{ij} denotes the Kronecker delta symbol. One also observes that

$$\left| D_i \varphi\left(\frac{x}{k}\right) \frac{x_j}{k} \right| \le C, \text{ for all } x \in \mathbb{R}^N, \ i, j = 1, 2, \cdots.$$
(2.10)

Denoting $f(s) = \lambda |s|^{p-2}s + |u|^{p-2}u \log |s|^p + \mu |s|^{q-2}s$ for all $s \in \mathbb{R}$, by means of (A.1) below, it holds that

$$\sum_{i,j=1}^{N} \int_{\mathbb{R}^{N}} D_{i}\varphi\left(\frac{x}{k}\right) \frac{x_{j}}{k} D_{\xi_{i}}\mathcal{L}(x,u,\nabla u) D_{j}udx + \int_{\mathbb{R}^{N}} \varphi\left(\frac{x}{k}\right) D_{\xi}\mathcal{L}(x,u,\nabla u) \cdot \nabla udx$$
$$- \int_{\mathbb{R}^{N}} \left[D\varphi\left(\frac{x}{k}\right) \frac{x}{k} \mathcal{L}(x,u,\nabla u) + N\varphi\left(\frac{x}{k}\right) \mathcal{L}(x,u,\nabla u) \right] dx$$
$$= \int_{\mathbb{R}^{N}} \left[\varphi\left(\frac{x}{k}\right) x \cdot \nabla u \right] f(u) dx.$$

Thanks to (2.10), $\varphi(\frac{x}{k}) \to 1$ and $\nabla \varphi(\frac{x}{k}) \cdot \frac{x}{k} \to 0$ as $k \to +\infty$, we obtain

$$\sum_{i,j=1}^{N} \int_{\mathbb{R}^{N}} D_{i}\varphi\left(\frac{x}{k}\right) \frac{x_{j}}{k} D_{\xi_{i}}\mathcal{L}(x,u,\nabla u) D_{j}udx + \int_{\mathbb{R}^{N}} \varphi\left(\frac{x}{k}\right) D_{\xi}\mathcal{L}(x,u,\nabla u) \cdot \nabla udx$$
$$- \int_{\mathbb{R}^{N}} \left[D\varphi\left(\frac{x}{k}\right) \frac{x}{k} \mathcal{L}(x,u,\nabla u) + N\varphi\left(\frac{x}{k}\right) \mathcal{L}(x,u,\nabla u) \right] dx$$
$$\rightarrow \int_{\mathbb{R}^{N}} |\nabla u|^{p} dx - N \int_{\mathbb{R}^{N}} \frac{1}{p} |\nabla u|^{p} dx = -\frac{N-p}{p} \int_{\mathbb{R}^{N}} |\nabla u|^{p} dx$$

as $k \to +\infty$. On the other hand, since $F(u) \in L^1(\mathbb{R}^N)$ for all $x \in X$ by (1.9), we shall exploit an integration by parts and the Lebesgue's Dominated Convergence theorem to reach

$$\int_{\mathbb{R}^{N}} \left[\varphi\left(\frac{x}{k}\right) x \cdot \nabla u \right] f(u) dx = -N \int_{\mathbb{R}^{N}} F(u) \varphi\left(\frac{x}{k}\right) dx - \int_{\mathbb{R}^{N}} \left[\nabla \varphi\left(\frac{x}{k}\right) \cdot \frac{x}{k} \right] F(u) dx$$
$$\rightarrow -N \int_{\mathbb{R}^{N}} F(u) dx$$

as $k \to +\infty$. So, we can conclude the equality

Journal of Differential Equations 421 (2025) 1-49

$$\frac{N-p}{p} \int_{\mathbb{R}^N} |\nabla u|^p dx = N \int_{\mathbb{R}^N} \left(\frac{\lambda - 1}{p} |u|^p + \frac{1}{p} |u|^p \log |u|^p + \frac{\mu}{q} |s|^q \right) dx.$$
(2.11)

Multiplying the nontrivial solution $u \in X$ on both sides of Eq. (2.3), one has that

$$\int_{\mathbb{R}^N} |\nabla u|^p dx = \int_{\mathbb{R}^N} (\lambda |u|^p + |u|^p \log |u|^p + \mu |u|^q) dx.$$
(2.12)

By multiplying $\frac{N}{n}$ in (2.12) and then minus (2.11), we get the desired identity (2.4).

Step 3. If the nontrivial solution $u(x) \ge 0$ for all $x \in \mathbb{R}^N$, then u(x) > 0 for all $x \in \mathbb{R}^N$.

Choosing a sufficiently small $\epsilon > 0$, we have

$$\Delta_p u = -\lambda u^{p-1} - u^{p-1} \log u^p - \mu u^{q-1} \le \xi(u) \text{ in } \{ x \in \mathbb{R}^N : 0 < u(x) < \epsilon \},\$$

where $\xi(0) \triangleq \lim_{s \to 0^+} \xi(s) = 0$ and $\xi : (0, +\infty) \to \mathbb{R}$ is defined by

$$\xi(s) = \begin{cases} -\lambda s^{p-1} - s^{p-1} \log s^p, & \text{if } \mu > 0, \\ -(\lambda + \mu)s^{p-1} + s^{p-1} \log s^p, & \text{if } \mu \le 0. \end{cases}$$

Clearly, ξ is continuous and nondecreasing when s > 0 is small enough. It is simple to calculate that $\xi(\sqrt[p]{e^{-\lambda}}) = 0$ if $\mu > 0$, and $\xi(\sqrt[p]{e^{-(\lambda+\mu)}}) = 0$ if $\mu \le 0$. Since $u(x) \ge 0$ for all $x \in \mathbb{R}^N$, then we apply the Step 1 and [52, Theorem 5] to finish the proof. \Box

3. The semiclassical problem

In this section, we shall contemplate the existence and concentration behavior of positive normalized solutions for a class of *p*-Laplacian equations with logarithmic nonlinearities. Nevertheless, first of all, let us consider the existence of positive solutions to the problem

$$\begin{cases} -\Delta_{p}u + \mu|u|^{p-2}u = \lambda|u|^{p-2}u + |u|^{p-2}u\log|u|^{p}, \text{ in } \mathbb{R}^{N}, \\ \int_{\mathbb{R}^{N}} |u|^{p}dx = a^{p}, \end{cases}$$
(3.1)

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2}\nabla u)$ denotes the usual *p*-Laplacian operator with $2 \le p < N$, $\mu \in [-1, +\infty)$ is a fixed constant and $\lambda \in \mathbb{R}$ is known as the Lagrange multiplier.

In general, to solve Problem (3.1), we look for critical points of the following variational functional

$$I_{\mu}(u) = \frac{1}{p} \int_{\mathbb{R}^{N}} \left[|\nabla u|^{p} + (\mu + 1)|u|^{p} \right] dx + \int_{\mathbb{R}^{N}} F_{1}(u)v dx - \int_{\mathbb{R}^{N}} F_{2}(u) dx$$

restricted to the sphere S(a) defined by

$$S(a) = \left\{ u \in X : \int_{\mathbb{R}^N} |u|^p dx = a^p \right\}.$$

Recalling Lemma 2.2, it follows that $(X, \|\cdot\|)$ is a reflexive and separable Banach space. Additionally, note that the imbedding $X \hookrightarrow W^{1,p}(\mathbb{R}^N)$ and $X \hookrightarrow L^{F_1}(\mathbb{R}^N)$ are continuous. As a consequence, we are derived from Section 2 that $I_{\mu} \in C^1(X, \mathbb{R})$ with

$$I'_{\mu}(u)v = \int_{\mathbb{R}^N} \left[|\nabla u|^{p-2} \nabla u \nabla v + (\mu+1)|u|^{p-2}v \right] dx + \int_{\mathbb{R}^N} F'_1(u)v dx - \int_{\mathbb{R}^N} F'_2(u)v dx, \forall v \in X.$$

Next, we will prove the following result for Problem (3.1).

Theorem 3.1. Let $2 \le p < N$. Then, there is a constant $\tilde{a} = \tilde{a}(\mu) > 0$ such that Problem (3.1) has a couple solution $(u, \lambda) \in X \times \mathbb{R}$ for all $a > \tilde{a}$, where u(x) > 0 for all $x \in \mathbb{R}^N$ and $\lambda < 0$.

The proof of the above theorem will be divided into several lemmas.

Lemma 3.2. Let $2 \le p < N$, the functional I_{μ} is coercive and bounded from below on S(a).

Proof. In view of the property-(\mathbf{P}_2) in Section 2, for every fixed $\tilde{q} \in \left(p, \frac{p^2}{N}\right)$, there exists a constant $C_{\tilde{q}} > 0$ such that

$$|F_2'(s)| \le C_{\tilde{q}}|s|^{\tilde{q}-1}, \ \forall s \in \mathbb{R}.$$

Moreover, by the Gagliardo-Nirenberg inequality (1.12),

$$\begin{split} I_{\mu}(u) &= \frac{1}{p} \int_{\mathbb{R}^{N}} \left[|\nabla u|^{p} + (\mu+1)|u|^{p} \right] dx - \frac{1}{p} \int_{\mathbb{R}^{N}} |u|^{p} \log |u|^{p} dx \\ &\geq \frac{1}{p} \int_{\mathbb{R}^{N}} |\nabla u|^{p} dx + \int_{\mathbb{R}^{N}} F_{1}(u) dx - C_{q} \mathbb{C}_{N, p, \tilde{q}} a^{\tilde{q}(1-\beta_{\tilde{q}})} \left(\int_{\mathbb{R}^{N}} |\nabla u|^{p} dx \right)^{\frac{\tilde{q}\beta_{\tilde{q}}}{p}}. \end{split}$$

Since $\tilde{q} \in \left(p, \frac{p^2}{N}\right)$, then $\tilde{q}\beta_{\tilde{q}} < p$ by (1.13). Moreover, adopting (2.2), we see that $\int_{\mathbb{R}^N} F_1(u)dx \to +\infty$ as $\|u\|_{F_1} \to \infty$. These facts reveal the proof of this lemma. \Box

As a direct consequence of Lemma 3.2, the real number

$$\mathcal{I}_{\mu,a} = \inf_{u \in S(a)} I_{\mu}(u)$$

is well-defined. Then, we are going to establish some properties of $\mathcal{I}_{\mu,a}$ with respect to the parameter $\mu \in [-1, +\infty)$.

Lemma 3.3. Let $2 \le p < N$, then there exists a constant $\tilde{a} = \tilde{a}(\mu) > 0$ such that $\mathcal{I}_{\mu,a} < 0$ for all $a > \tilde{a}$ and $\mu \in [-1, +\infty)$.

Proof. Given some fixed $\psi \in X \setminus \{0\}$ and t > 0, it follows from some simple calculations that

$$I_{\mu}(t\psi) = \frac{t^{p}}{p} \int_{\mathbb{R}^{N}} \left[|\nabla\psi|^{p} + (\mu+1)|\psi|^{p} \right] dx - \frac{t^{p}}{p} \int_{\mathbb{R}^{N}} |\psi|^{p} \log|\psi|^{p} dx - t^{p} \log t \int_{\mathbb{R}^{N}} |\psi|^{p} dx$$
$$\to -\infty$$

as $t \to +\infty$. Hence, there is a sufficiently large constant $\tilde{t} > 0$ such that

$$I_{\mu}(t\psi) \leq -1$$
 for all $t > \tilde{t}$.

Then, we can choose $\tilde{a} = \tilde{t} |\psi|_p$ to reach the statement. \Box

Lemma 3.4. Let $2 \le p < N$. Fix $\mu \in [-1, +\infty)$ and let $0 < a_1 < a_2 < +\infty$, then $\frac{a_1^{\mu}}{a_2^{\mu}} \mathcal{I}_{\mu,a_2} < \mathcal{I}_{\mu,a_1}$.

Proof. Since $I_{\mu}(u) = I_{\mu}(|u|)$ for each $u \in X$, without loss of generality, we suppose that $\{u_n\} \subset S(a_1)$ is a nonnegative minimizing sequence with respect to \mathcal{I}_{μ,a_1} , that is,

$$I_{\mu}(u_n) \to \mathcal{I}_{\mu,a_1}, \text{ as } n \to +\infty.$$

Choosing $v_n = \xi u_n$, then $v_n \in S(a_2)$ for every $n \in \mathbb{N}$, where $\xi \triangleq \frac{a_2}{a_1} > 1$. It follows from some simple calculations that

$$\mathcal{I}_{\mu,a_2} \le I_{\mu}(v_n) = \xi^p I_{\mu}(u_n) - \frac{1}{p} \xi^p \log \xi^p \int_{\mathbb{R}^N} |u_n|^p dx = \xi^p I_{\mu}(u_n) - \frac{1}{p} a_1^p \xi^p \log \xi^p.$$

Letting $n \to +\infty$ and using the fact that $\xi > 1$, there holds

$$\mathcal{I}_{\mu,a_2} \leq \xi^p \mathcal{I}_{\mu,a_1} - \frac{1}{p} a_1^p \xi^p \log \xi^p < \xi^p \mathcal{I}_{\mu,a_1}$$

that is,

$$\frac{a_1^p}{a_2^p}\mathcal{I}_{\mu,a_2} < \mathcal{I}_{\mu,a_1},$$

finishing the proof of this lemma. \Box

Borrowing some ideas from [8, Theorem 3.2], we derive the compactness theorem on S(a) which plays a pivotal role in the proof of Theorem 3.1.

Theorem 3.5. (Compactness theorem on S(a)) Let $2 \le p < N$. Suppose that $a > \tilde{a}$ and $\{u_n\} \subset S(a)$ is a minimizing sequence with respect to $\mathcal{I}_{\mu,a}$, then, for some subsequence either i) $\{u_n\}$ is strongly convergent in X,

ii) There exists $\{y_n\} \subset \mathbb{R}^N$ such that the sequence $v_n(x) = u_n(x + y_n)$ is strongly convergent to a function $v \in S(a)$ in X with $I_{\mu}(v) = \mathcal{I}_{\mu,a}$, where $|y_n| \to +\infty$ along a subsequence.

Proof. Since I_{μ} is coercive on S(a) by Lemma 3.2, the sequence $\{u_n\}$ is bounded, and then, $u_n \rightarrow u$ in X for some subsequence. If $u \neq 0$ and $|u|_p = b \neq a$, we must have $b \in (0, a)$. By the Brézis-Lieb Lemma (see e.g. [58]),

$$|u_n|_p^p = |u_n - u|_p^p + |u|_p^p + o_n(1).$$

Setting $v_n = u_n - u$, $d_n = |v_n|_p$ and supposing that $|v_n|_p \to d$, we get $a^p = b^p + d^p$. From $d_n \in (0, a)$ for *n* large enough, we apply Lemma 2.4 to see that

$$\mathcal{I}_{\mu,a} + o_n(1) = I_{\mu}(u_n) = I_{\mu}(v_n) + I_{\mu}(u) + o_n(1) \ge \mathcal{I}_{\mu,d_n} + \mathcal{I}_{\mu,b} + o_n(1).$$

Thereby, by Lemma 3.4,

$$\mathcal{I}_{\mu,a} + o_n(1) \ge \frac{d_n^p}{a^p} \mathcal{I}_{\mu,a} + \mathcal{I}_{\mu,b} + o_n(1).$$

Letting $n \to +\infty$, we find

$$\mathcal{I}_{\mu,a} \ge \frac{d^p}{a^p} \mathcal{I}_{\mu,a} + \mathcal{I}_{\mu,b}.$$
(3.2)

Since $b \in (0, a)$, using again Lemma 3.4 in (3.2), we get the following inequality

$$\mathcal{I}_{\mu,a} > \frac{d^p}{a^p} \mathcal{I}_{\mu,a} + \frac{b^p}{a^p} \mathcal{I}_{\mu,b} = \left(\frac{d^p}{a^p} + \frac{b^p}{a^p}\right) \mathcal{I}_{\mu,a} = \mathcal{I}_{\mu,a},$$

which is absurd. This asserts that $|u|_p = a$, or equivalently, $u \in S(a)$. As $|u_n|_p = |u|_p = a$, $u_n \rightarrow u$ in $L^p(\mathbb{R}^N)$ and $L^p(\mathbb{R}^N)$ is reflexive, it is well-known that

$$u_n \to u \text{ in } L^p(\mathbb{R}^N). \tag{3.3}$$

This combined with interpolation theorem in the Lebesgue space and property- (\mathbf{P}_2) reveal

$$\int_{\mathbb{R}^N} F_2(u_n) dx \to \int_{\mathbb{R}^N} F_2(u) dx.$$
(3.4)

These limits together with $\mathcal{I}_{\mu,a} = \lim_{n \to +\infty} I_{\mu}(u_n)$ and $F_1 \ge 0$ in property-(\mathbf{P}_1) indicate that

$$\mathcal{I}_{\mu,a} \geq I_{\mu}(u).$$

As $u \in S(a)$, therefore $I_{\mu}(u) = \mathcal{I}_{\mu,a}$, then

$$\lim_{n \to +\infty} I_{\mu}(u_n) = I_{\mu}(u),$$

that combines with (3.3) and (3.4) to give

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} |\nabla u_n|^p dx = \int_{\mathbb{R}^N} |\nabla u|^p dx$$

and

$$\lim_{n\to\infty}\int_{\mathbb{R}^N}F_1(u_n)dx=\int_{\mathbb{R}^N}F_1(u)dx.$$

Recalling $F_1 \in (\Delta_2)$ by Lemma 2.2, jointly with the above two limits as well as (3.3), it enables to see that $u_n \to u$ in X.

Now, assume that u = 0, that is, $u_n \rightarrow 0$ in X. We claim that there exists C > 0 such that

$$\int_{\mathbb{R}^N} F_2(u_n) dx \ge C, \text{ for } n \in \mathbb{N} \text{ large enough.}$$
(3.5)

Otherwise, there is a subsequence of $\{u_n\}$, still denoted by itself, such that

$$\int_{\mathbb{R}^N} F_2(u_n) dx \to 0$$

as $n \to \infty$. It follows from Lemma 3.3 and (3.5) that

$$0 > \mathcal{I}_{\mu,a} = \lim_{n \to \infty} I_{\mu}(u_n) = \lim_{n \to \infty} \left(\frac{1}{p} \int_{\mathbb{R}^N} \left[|\nabla u_n|^p + (\mu+1)|u_n|^p \right] dx + \int_{\mathbb{R}^N} F_1(u_n) dx \right) \ge 0$$

which is impossible.

So, there are R, C > 0 and $\{y_n\} \subset \mathbb{R}^N$ such that

$$\int_{B_R(y_n)} |u_n|^p dx \ge C, \text{ for all } n \in \mathbb{N}.$$
(3.6)

If it is not the case, then we derive $u_n \to 0$ for all $p < s < p^*$ by the Vanishing lemma which yields that $F_2(u_n) \to 0$ in $L^1(\mathbb{R}^N)$ by property-(\mathbf{P}_2), a contradiction to (3.5). Recalling u = 0, we further have that $\{y_n\}$ is unbounded in \mathbb{R}^N . Define $v_n(x) = u_n(x + y_n)$, then $\{v_n\} \subset S(a)$ and it is also a minimizing sequence with respect to $\mathcal{I}_{\mu,a}$. Moreover, owing to (3.6), passing to a subsequence if necessary, there is a $v \in X \setminus \{0\}$ such that

$$v_n \rightarrow v$$
 in X and $v_n \rightarrow v$ a.e. in \mathbb{R}^N .

Repeating the calculations in the first part of the proof, it must conclude that $v_n \rightarrow v$ in X along a subsequence. The proof is completed. \Box

Proof of Theorem 3.1. By Lemma 3.2, there exists a bounded minimizing sequence $\{u_n\} \subset S(a)$ with respect to $\mathcal{I}_{\mu,a}$, that is, $I_{\mu}(u_n) \to \mathcal{I}_{\mu,a}$. Thanks to Theorem 3.5, there exists a $u_a \in S(a)$ with $I_{\mu}(u_a) = \mathcal{I}_{\mu,a}$. Therefore, by the Lagrange multiplier theorem, there exists $\lambda_a \in \mathbb{R}$ such that

$$I'_{\mu}(u_a) = \lambda_a \Psi'(u_a) \text{ in } X^*, \tag{3.7}$$

where $\Psi: X \to \mathbb{R}$ is given by

$$\Psi(u) = \frac{1}{p} \int_{\mathbb{B}^N} |u|^p dx, \ u \in X.$$

Thereby, according to (3.7), the couple $(u_a, \lambda_a) \in S(a) \times \mathbb{R}$ satisfies the following equation

$$-\Delta_p u + \mu |u|^{p-2} u = \lambda |u|^{p-2} u + |u|^{p-2} u \log |u|^p, \text{ in } \mathbb{R}^N.$$

Let $u_a \in S(a)$ be a test function on the both sides for the above equation, it holds that

$$\int_{\mathbb{R}^N} \left(|\nabla u_a|^p + \mu |u_a|^p \right) dx = \lambda_a a^p + \int_{\mathbb{R}^N} |u_a|^p \log |u_a|^p dx$$

which indicates that

$$\mathcal{I}_{\mu,a} = I_{\mu}(u_a) = \frac{1}{p} \int_{\mathbb{R}^N} |u_a|^p dx + \frac{\lambda_a}{p} a^p \ge \frac{\lambda_a}{p} a^p.$$

Due to Lemma 3.3, one sees that $\lambda_a < 0$. Since $u \in S(a)$ implies that $|u| \in S(a)$ and $I_{\mu}(u) = I_{\mu}(|u|)$ for all $u \in X$ which give that

$$|u_a| \in S(a)$$
 and $\mathcal{I}_{\mu,a} = I_{\mu}(u_a) \ge I_{\mu}(|u_a|) \ge \mathcal{I}_{\mu,a}$.

So, we can replace u_a with $|u_a|$ and then, without loss of generality, we shall suppose that $u_a \ge 0$. A very similar arguments in Step 3 in the proof of Theorem 2.5 show that u_a is positive. The proof is completed. \Box

Thanks to Theorem 3.1, we immediately have the following result whose detailed proof is omitted.

Corollary 3.6. Let $2 \le p < N$. If $a > \tilde{a}$ and $-1 \le \mu_1 < \mu_2 < +\infty$ are fixed, then $\mathcal{I}_{\mu_1,a} < \mathcal{I}_{\mu_2,a} < 0$.

From now on, we begin investigating the existence and concentration behavior of positive solutions for (1.1)-(1.2). To the aims, we consider the variational functional $I_{\varepsilon}: X \to \mathbb{R}$ given by

$$I_{\varepsilon}(u) = \frac{1}{p} \int_{\mathbb{R}^N} \left[|\nabla u|^p + (V(\varepsilon x) + 1)|u|^p \right] dx + \int_{\mathbb{R}^N} F_1(u) dx - \int_{\mathbb{R}^N} F_2(u) dx$$
(3.8)

restricted to the sphere S(a) and the minimization problem

$$\mathcal{I}_{\varepsilon,a} = \inf_{u \in S(a)} I_{\varepsilon}(u).$$

According to Theorem 3.5, it is significant to derive a similar compactness theorem for I_{ε} on S(a). So, we shall focus on verifying it. Let us introduce the two functionals $I_0, I_{\infty} : S(a) \to \mathbb{R}$ defined by

$$\begin{cases} I_{0}(u) = \frac{1}{p} \int_{\mathbb{R}^{N}} \left[|\nabla u|^{p} + (V_{0} + 1)|u|^{p} \right] dx + \int_{\mathbb{R}^{N}} F_{1}(u) dx - \int_{\mathbb{R}^{N}} F_{2}(u) dx, \\ I_{\infty}(u) = \frac{1}{p} \int_{\mathbb{R}^{N}} \left[|\nabla u|^{p} + (V_{\infty} + 1)|u|^{p} \right] dx + \int_{\mathbb{R}^{N}} F_{1}(u) dx - \int_{\mathbb{R}^{N}} F_{2}(u) dx \end{cases}$$

The corresponding minimization problems are given by

$$\mathcal{I}_{0,a} = \inf_{u \in S(a)} I_0(u) \text{ and } \mathcal{I}_{\infty,a} = \inf_{u \in S(a)} I_\infty(u).$$

Since $-1 \le V_0 < V_{\infty} < +\infty$ by (\hat{V}_1) , we are derived from Corollary 3.6 that

$$\mathcal{I}_{0,a} < \mathcal{I}_{\infty,a} < 0 \text{ for all } a > \tilde{a} > 0.$$
(3.9)

Lemma 3.7. Let $2 \le p < N$ and $a > \tilde{a} > 0$, then $\limsup_{\varepsilon \to 0^+} \mathcal{I}_{\varepsilon,a} \le \mathcal{I}_{0,a}$. In particular, there exists a sufficiently small $\varepsilon^* > 0$ such that $\mathcal{I}_{\varepsilon,a} < \mathcal{I}_{\infty,a}$ for all $\varepsilon \in (0, \varepsilon^*)$.

Proof. Adopting Theorem 3.1, for all $a > \tilde{a} > 0$, there is a $u_0 \in S(a)$ such that $I_0(u_0) = \mathcal{I}_{0,a}$. So,

$$\mathcal{I}_{\varepsilon,a} \leq I_{\varepsilon}(u_0) = \frac{1}{p} \int_{\mathbb{R}^N} \left[|\nabla u_0|^p + (V(\varepsilon x) + 1)|u_0|^p \right] dx + \int_{\mathbb{R}^N} F_1(u_0) dx - \int_{\mathbb{R}^N} F_2(u_0) dx.$$

Applying the Lebesgue's theorem and taking the limit as $\varepsilon \to 0^+$, there holds

$$\limsup_{\varepsilon \to 0^+} \mathcal{I}_{\varepsilon,a} \le \limsup_{\varepsilon \to 0^+} I_{\varepsilon}(u_0) = I_0(u_0) = \mathcal{I}_{0,a}$$

finishing the first part of the lemma. Due to (3.9), one could find such a constant $\varepsilon^* > 0$ such that $\mathcal{I}_{\varepsilon,a} < \mathcal{I}_{\infty,a}$ for all $\varepsilon \in (0, \varepsilon^*)$. The proof is completed. \Box

Lemma 3.8. Let $2 \le p < N$ and $a > \tilde{a} > 0$. If $\varepsilon \in (0, \varepsilon^*)$ is fixed and suppose that $\{u_n\} \subset S(a)$ such that $I_{\varepsilon}(u_n) \rightarrow \hat{d} < \frac{\mathcal{I}_{0,a} + \mathcal{I}_{\infty,a}}{2}$, then there is a $u \ne 0$ such that $u_n \rightharpoonup u$ in X along a subsequence.

Proof. Arguing as Lemma 3.2, one deduces that $\{u_n\}$ is bounded in X. Passing to a subsequence if necessary, there is a $u \in X$ such that $u_n \rightarrow u$ in X and $u_n \rightarrow u$ a.e. in \mathbb{R}^N . To prove that $u \neq 0$, let us suppose it by a contradiction and assume that $u \equiv 0$. Hence,

$$\hat{d} + o_n(1) = I_{\varepsilon}(u_n) = I_{\infty}(u_n) + \frac{1}{p} \int_{\mathbb{R}^N} [V(\varepsilon x) - V_{\infty}] |u_n|^p dx.$$

Due to (V_2), given an arbitrary $\epsilon > 0$, there is an R > 0 such that

$$V(x) \ge V_{\infty} - \epsilon$$
, for all $|x| \ge R$

which indicates that

$$\hat{d} + o_n(1) \ge I_{\infty}(u_n) + \frac{1}{p} \int_{B_{R/\varepsilon}(0)} [V(\varepsilon x) - V_{\infty}] |u_n|^p dx - \frac{\epsilon}{p} \int_{B_{R/\varepsilon}^c(0)} |u_n|^p dx.$$

Since $\{u_n\}$ is bounded in X and $u_n \to 0$ in $L^p(B_{R/\varepsilon}(0))$, one has that

$$\hat{d} + o_n(1) \ge I_{\infty}(u_n) - C\epsilon \ge \mathcal{I}_{\infty,a} - C\epsilon$$

for some C > 0 independent of ϵ . Let us tend $\epsilon \to 0^+$, then $\hat{d} \ge \mathcal{I}_{\infty,a}$ which is impossible due to the facts that $\hat{d} < \frac{\mathcal{I}_{0,a} + \mathcal{I}_{\infty,a}}{2}$ and (3.9). So, the proof of this lemma concludes. \Box

Lemma 3.9. Let $2 \le p < N$ and $a > \tilde{a} > 0$. Suppose $\varepsilon \in (0, \varepsilon^*)$ to be fixed and let $\{u_n\} \subset X$ be $a (PS)_{\hat{d}}$ sequence for I_{ε} constrained to S(a) with $\hat{d} < \frac{\mathcal{I}_{0,a} + \mathcal{I}_{\infty,a}}{2}$, namely

$$I_{\varepsilon}(u_n) \to \hat{d} \text{ and } \|I'_{\varepsilon}|_{S(a)}\|_{X^*} \to 0 \text{ as } n \to \infty,$$

then, up to a subsequence if necessary, there is a $u \in X$ such that $u_n \rightharpoonup u$ in X. Moreover if $u_n \not\rightarrow u$ in X, then there exists a $\hat{\delta} > 0$ independent of $\varepsilon \in (0, \varepsilon^*)$ such that, by decreasing ε^* if necessary, there holds

$$\liminf_{n\to\infty}\int_{\mathbb{R}^N}|v_n|^pdx\geq\hat{\delta}.$$

Proof. We define the functional $\Psi : X \to \mathbb{R}$ by

$$\Psi(u) = \frac{1}{p} \int_{\mathbb{B}^N} |u|^p dx, \ u \in X,$$

it follows that $S(a) = \Psi^{-1}(\{a^p/p\})$. Hence, adopting [58, Proposition 5.12], there exists $\{\lambda_n\} \subset \mathbb{R}$ such that

$$||I_{\varepsilon}'(u_n) - \lambda_n \Psi'(u_n)||_{X^*} \to 0 \text{ as } n \to \infty.$$

Since $\{u_n\}$ is bounded in X, we easily get that $\{\lambda_n\}$ is bounded in \mathbb{R} . Passing to a subsequence if necessary, there is a $\lambda \in \mathbb{R}$ that may depend on ε such that $\lambda_n \to \lambda$ and so

$$||I_{\varepsilon}'(u_n) - \lambda \Psi'(u_n)||_{X^*} \to 0 \text{ as } n \to \infty$$

which immediately shows us that

$$I_{\varepsilon}'(u) - \lambda \Psi'(u) = 0$$
 in X^* .

Combining Lemma 2.4 and the Brézis-Lieb lemma, one has that

$$I_{\varepsilon}'(v_n)v_n - \lambda \Psi'(v_n)v_n = I_{\varepsilon}'(u_n)u_n - I_{\varepsilon}'(u_n)u - \lambda \Psi'(u_n)u_n + \lambda \Psi'(u_n)u + o_n(1)$$
$$= I_{\varepsilon}'(u_n)u_n - \lambda_n \Psi'(u_n)u_n - I_{\varepsilon}'(u_n)u + \lambda \Psi'(u_n)u + o_n(1)$$
$$= o_n(1)$$

jointly with $F'_1(s)s \ge 0$ for all $s \in \mathbb{R}$ in property- (\mathbf{P}_1) and property- (\mathbf{P}_2) implies that

$$\int_{\mathbb{R}^N} [|\nabla v_n|^p + (V(\varepsilon x) + 1 - \lambda)|v_n|^p] dx \le C_{\tilde{q}} \int_{\mathbb{R}^N} |v_n|^{\tilde{q}} dx + o_n(1)$$

for some $\tilde{q} \in (p, p^*)$. We claim that there is a $\lambda^* < 0$ independent of $\varepsilon \in (0, \varepsilon^*)$ such that

$$\lambda \leq \lambda^*, \ \forall \varepsilon \in (0, \varepsilon^*).$$

Indeed, due to $\{u_n\} \subset S(a)$, we find that

$$\hat{d} = \lim_{n \to \infty} I_{\varepsilon}(u_n) = \lim_{n \to \infty} \left[I_{\varepsilon}(u_n) - \frac{1}{p} \left(I_{\varepsilon}'(u_n)u_n - \lambda \Psi'(u_n)u_n \right) \right] \ge \frac{\lambda}{p} a^p$$

showing the claim. As a consequence, owing to $V_0 \ge -1$ by (\hat{V}_1) , there holds

$$\int_{\mathbb{R}^N} \left(|\nabla v_n|^p - \lambda^* |v_n|^p \right) dx \le C_{\tilde{q}} \int_{\mathbb{R}^N} |v_n|^{\tilde{q}} dx + o_n(1), \ \forall \varepsilon \in (0, \varepsilon^*).$$

It follows from the continuous imbedding $W^{1,p}(\mathbb{R}^N) \hookrightarrow L^q(\mathbb{R}^N)$ with $q \in (p, p^*)$ that

$$\|v_n\|_{W^{1,p}(\mathbb{R}^N)}^p \le C_1 |v_n|_{\tilde{q}}^{\tilde{q}} + o_n(1) \le C_2 \|v_n\|_{W^{1,p}(\mathbb{R}^N)}^{\tilde{q}}, \ \forall \varepsilon \in (0, \varepsilon^*),$$

where $C_1, C_2 > 0$ are independent of $\varepsilon \in (0, \varepsilon^*)$. Because $v_n \neq 0$ in X and $\tilde{q} > p$, without loss of generality, we can suppose that

$$\liminf_{n \to \infty} \|v_n\|_{W^{1,p}(\mathbb{R}^N)} \ge C_3, \ \forall \varepsilon \in (0, \varepsilon^*),$$
(3.10)

where $C_3 > 0$ is independent of $\varepsilon \in (0, \varepsilon^*)$. Otherwise, if $v_n \to 0$ in $W^{1,p}(\mathbb{R}^N)$, then $F'_2(v_n)v_n \to 0$ in $L^1(\mathbb{R}^N)$ by property- (\mathbf{P}_2) which together with $I'_{\varepsilon}(v_n)v_n - \lambda \Psi'(v_n)v_n = o_n(1)$ and $F'_2(s)s \ge 0$ for all $s \in \mathbb{R}$ by property- (\mathbf{P}_1) yields that $F_1(v_n) \to 0$ in $L^1(\mathbb{R}^N)$. Recalling Lemma 2.2, we can conclude that $v_n \to 0$ in $L^{F_1}(\mathbb{R}^N)$ and so $v_n \to 0$ in X, a contradiction and (3.10) follows. Hence,

$$\liminf_{n\to\infty} |v_n|_{\tilde{q}}^{\tilde{q}} \ge C_2^{-1}C_3^p, \ \forall \varepsilon \in (0,\varepsilon^*).$$

Adopting (1.12) and since $|\nabla v_n|_p^p$ is bounded, we can derive the proof of this lemma. \Box

Theorem 3.10. Let $2 \le p < N$ and $a > \tilde{a} > 0$. If $\varepsilon \in (0, \varepsilon^*)$ is fixed, then the functional $I_{\varepsilon}|_{S(a)}$ satisfies the $(PS)_{\hat{d}}$ condition with $\hat{d} < \mathcal{I}_{0,a} + \Upsilon$, where $0 < \Upsilon \le \min\{\frac{1}{2}, \frac{\hat{\delta}}{a^p}\}(\mathcal{I}_{\infty,a} - \mathcal{I}_{0,a})$ and $\hat{\delta} > 0$ is determined by Lemma 3.9.

Proof. Let $\{u_n\} \subset X$ be a $(PS)_d$ sequence for $I_{\varepsilon}|_{S(a)}$ and define the functional $\Psi: X \to \mathbb{R}$ by

$$\Psi(u) = \frac{1}{p} \int_{\mathbb{B}^N} |u|^p dx, \ u \in X,$$

which reveals that $S(a) = \Psi^{-1}(\{a^p/p\})$. Thus, adopting [58, Proposition 5.12], there exists $\{\lambda_n\} \subset \mathbb{R}$ such that

$$||I_{\varepsilon}'(u_n) - \lambda_n \Psi'(u_n)||_{X^*} \to 0 \text{ as } n \to \infty.$$

According to Lemma 3.3, $\{u_n\}$ is bounded in X and then there is a $u \in X$ such that $u_n \rightharpoonup u$ in X and $u_n \rightarrow u$ a.e. in \mathbb{R}^N along a subsequence. Denoting $v_n \triangleq u_n - u$, if $v_n \not\rightarrow 0$ in X, then Lemma 3.9 ensures that

$$\liminf_{n \to \infty} \int_{\mathbb{R}^N} |v_n|^p dx \ge \hat{\delta}.$$
(3.11)

Let $d_n = |v_n|_p$ and $|u|_p = b$, so we can assume that $|v_n|_p \to d > 0$ and b > 0 by (3.11) and Lemma 3.8, respectively. Moreover, it holds that $a^p = b^p + d^p$ by the Brézis-Lieb lemma. Via exploiting a similar argument explored in Lemma 3.8, by means of $v_n \to 0$ in X, one can show that $I_{\varepsilon}(v_n) \ge \mathcal{I}_{\infty,d_n} + o_n(1)$ which together with Lemma 2.4 and $V(x) \ge V_0$ for all $x \in \mathbb{R}^N$ by (\hat{V}_1) gives that

$$\hat{d} + o_n(1) = I_{\varepsilon}(u_n) = I_{\varepsilon}(v_n) + I_{\varepsilon}(u) + o_n(1) \ge \mathcal{I}_{\infty,d_n} + \mathcal{I}_{0,b} + o_n(1)$$

Since $d_n \in (0, a)$ for $n \in \mathbb{N}$ large enough and $b \in (0, a)$, we argue as Lemma 3.4 to derive

$$\hat{d} + o_n(1) \ge \frac{d_n^p}{a^p} \mathcal{I}_{\infty,a} + \frac{b^p}{a^p} \mathcal{I}_{0,a}$$

Letting $n \to \infty$ and recalling $a^p = b^p + d^p$, one has that

$$\Upsilon > \frac{\hat{\delta}}{a^p} (\mathcal{I}_{\infty,a} - \mathcal{I}_{0,a})$$

violating to the definition of Υ . So, we must conclude that $v_n \to 0$ in X, that is $u_n \to u$ in X. One further obtains that $u \in S(a)$ and

$$-\Delta_p u + V(\varepsilon x)|v|^{p-2}u = \lambda |u|^{p-2}v + |u|^{p-2}v \log |v|^p \text{ in } \mathbb{R}^N,$$

where $\lambda \leq \lambda^* < 0$ comes from Lemma 3.9. The proof is completed. \Box

In what follows, according to (V_2) , we fix some sufficiently small ρ_0 , $r_0 > 0$ to satisfy

- $\overline{B_{\rho_0}(x^i)} \cap \overline{B_{\rho_0}(x^j)} = \emptyset$ for $i \neq j$ and $i, j \in \{1, \dots, l\};$ • $\bigcup_{i=1}^{l} B_{\rho_0}(x^i) \subset B_{r_0}(0);$
- $K_{\frac{\rho_0}{2}} = \bigcup_{i=1}^{\ell} \overline{B_{\frac{\rho_0}{2}}(x^i)}.$

Define the function $Q_{\varepsilon}: X \setminus \{0\} \to \mathbb{R}$ by

$$Q_{\varepsilon}(u) = \frac{\int\limits_{\mathbb{R}^{N}} \chi(\varepsilon x) |u|^{p} dx}{\int\limits_{\mathbb{R}^{N}} |u|^{p} dx},$$

where $\chi : \mathbb{R}^N \to \mathbb{R}^N$ is given by

$$\chi(x) = \begin{cases} x, & \text{if } |x| \le r_0, \\ \frac{r_0 x}{|x|}, & \text{if } |x| > r_0. \end{cases}$$

With Theorem 3.10 in hands, we now focus on establishing the existence of (*PS*) sequences for the variational functional I_{ε} constrained on S(a).

Lemma 3.11. Let $2 \le p < N$ and $a > \tilde{a} > 0$. Suppose $\varepsilon \in (0, \varepsilon^*)$ to be fixed, decreasing $\varepsilon^* > 0$ if necessary, there is $\delta^* > 0$ such that if $u \in S(a)$ and $I_{\varepsilon}(u) \le I_{0,a} + \delta^*$, then

$$Q_{\varepsilon}(u) \in K_{\frac{\rho_0}{2}}, \ \forall \varepsilon \in (0, \varepsilon^*).$$

Proof. Suppose, by contradiction, that for all $n \in \mathbb{N}$, there exist $\varepsilon_n \to 0$ and $\{u_n\} \subset S(a)$ such that

$$\mathcal{I}_{0,a} \leq I_{\varepsilon_n}(u_n) \leq \mathcal{I}_{0,a} + \frac{1}{n},$$

and

$$Q_{\varepsilon_n}(u_n) \notin K_{\frac{\rho_0}{2}}.$$

Obviously, one has that

$$\mathcal{I}_{0,a} \le I_0(u_n) \le I_{\varepsilon_n}(u_n) \le \mathcal{I}_{0,a} + \frac{1}{n}$$

showing that $\{u_n\} \subset S(a)$ is a minimizing sequence with respect to $\mathcal{I}_{0,a}$. Thanks to Theorem 3.5, passing to a subsequence if necessary, one of the following alternatives holds true

i) There is a function $u \in S(a)$ such that $u_n \to u$ in X as $n \to \infty$;

ii) There exists a sequence $\{y_n\} \subset \mathbb{R}^N$ with $|y_n| \to +\infty$ such that $v_n = u_n(\cdot + y_n) \to v$ in X for some $v \in S(a)$.

We claim that i) cannot occur. Otherwise, adopting the definition of χ , one has

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} \chi(\varepsilon_n x) |u_n|^p dx = \lim_{n \to \infty} \int_{\mathbb{R}^N} \chi(0) |u|^p dx = 0.$$

From which, we conclude that $Q_{\varepsilon_n}(u) \in K_{\frac{\rho_0}{2}}$ for some sufficiently large $n \in \mathbb{N}$. It is impossible and so the claim follows. When ii) occurs, passing to a subsequence if necessary, we shall contemplate the following two cases:

ii)-(1). $|\varepsilon_n y_n| \to +\infty$ as $n \to \infty$. In this case, as a consequence of $v_n \to v$ in X, there holds

$$\begin{split} I_{\varepsilon_n}(u_n) &= \frac{1}{p} \int\limits_{\mathbb{R}^N} \left[|\nabla u_n|^p + (V(\varepsilon_n x) + 1)|u_n|^p \right] dx + \int\limits_{\mathbb{R}^N} F_1(u_n) dx - \int\limits_{\mathbb{R}^N} F_2(u_n) dx \\ &= \frac{1}{p} \int\limits_{\mathbb{R}^N} \left[|\nabla v_n|^p + (V(\varepsilon_n x + \varepsilon_n y_n) + 1)|v_n|^p \right] dx + \int\limits_{\mathbb{R}^N} F_1(v_n) dx - \int\limits_{\mathbb{R}^N} F_2(v_n) dx \\ &\to I_{\infty}(v). \end{split}$$

Since $I_{\varepsilon_n}(u_n) \leq \mathcal{I}_{0,a} + \frac{1}{n}$, we arrive at the inequality below

$$\mathcal{I}_{0,a} \ge I_{\infty}(v) \ge \mathcal{I}_{\infty,a}$$

which contradicts with (3.9).

ii)-(2). $\varepsilon_n y_n \to y$ for some $y \in \mathbb{R}^N$ as $n \to \infty$. In this case, a similar argument using the above calculations indicates that

$$\mathcal{I}_{V(y),a} \leq \mathcal{I}_{0,a}.$$

If $V(y) > V_0$, we follow a very similar approach explored in the proof of Corollary 3.6 to deduce that $\mathcal{I}_{V(y),a} > \mathcal{I}_{0,a}$ which is absurd. Thereby, $V(y) = V_0$ and $y = x^i$ for some $i \in \{0, 1, \dots, l\}$. Then one derives that

Journal of Differential Equations 421 (2025) 1-49

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} \chi(\varepsilon_n x) |u_n|^p dx = \lim_{n \to \infty} \int_{\mathbb{R}^N} \chi(\varepsilon_n x + \varepsilon_n y_n) |v_n|^p dx = x^i \int_{\mathbb{R}^N} |v|^p dx$$

which reveals that $\lim_{n \to \infty} Q_{\varepsilon_n}(u_n) = x^i \in K_{\frac{\rho_0}{2}}$. From which, we obtain that $Q_{\varepsilon_n}(u_n) \in K_{\frac{\rho_0}{2}}$ for some sufficiently large $n \in N$, a contradiction. The proof is completed. \Box

In the sequel, for $j \in \{1, \dots, l\}$, we need the following notations

•
$$\theta_{\varepsilon}^{j} = \{u \in S(a) : |Q_{\varepsilon}(u) - x^{j}| < \rho_{0}\},\$$

• $\partial \theta_{\varepsilon}^{j} = \{u \in S(a) : |Q_{\varepsilon}(u) - x^{j}| = \rho_{0}\},\$
• $\beta_{\varepsilon}^{j} = \inf_{u \in \theta_{\varepsilon}^{j}} I_{\varepsilon}(u) \text{ and } \tilde{\beta}_{\varepsilon}^{j} = \inf_{u \in \partial \theta_{\varepsilon}^{j}} I_{\varepsilon}(u).$

Lemma 3.12. Let $2 \le p < N$ and $a > \tilde{a} > 0$. Suppose $\varepsilon \in (0, \varepsilon^*)$ to be fixed, decreasing $\varepsilon^* > 0$ if necessary, for the constant $\Upsilon > 0$ in Theorem 3.10, there holds

$$\beta_{\varepsilon}^{j} < \mathcal{I}_{0,a} + \Upsilon \text{ and } \beta_{\varepsilon}^{j} < \tilde{\beta}_{\varepsilon}^{j}, \ \forall \varepsilon \in (0, \varepsilon^{*}).$$

Proof. According to Theorem 3.1, there is a function $u \in S(a)$ such that

$$I_0(u) = \mathcal{I}_{0,a}$$
 and $I'_0(u) = 0$ in X^* .

For $j \in \{1, \dots, l\}$, we define the function $\hat{u}_{\varepsilon}^{j} : \mathbb{R}^{N} \to \mathbb{R}$ by

$$\hat{u}_{\varepsilon}^{j} = u(x - x^{j}/\varepsilon).$$

By a simple change of variable, one has that

$$I_{\varepsilon}(\hat{u}_{\varepsilon}^{j}) = \frac{1}{p} \int_{\mathbb{R}^{N}} \left[|\nabla u|^{p} + (V(\varepsilon x + x^{j}) + 1)|u|^{p} \right] dx + \int_{\mathbb{R}^{N}} F_{1}(u) dx - \int_{\mathbb{R}^{N}} F_{2}(u) dx$$

which gives that

$$\limsup_{\varepsilon \to 0^+} I_{\varepsilon}(\hat{u}_{\varepsilon}^j) = I_0(u) = \mathcal{I}_{0,a}.$$
(3.12)

From which, decreasing $\varepsilon^* > 0$ if necessary,

$$I_{\varepsilon}(\hat{u}_{\varepsilon}^{j}) < \mathcal{I}_{0,a} + \frac{1}{4}\delta^{*}, \ \forall \varepsilon \in (0, \varepsilon^{*}),$$

where $\delta^* > 0$ comes from Lemma 3.11. Moreover, we easily deduce that $Q_{\varepsilon}(\hat{u}_{\varepsilon}^j) \to x^j$ as $\varepsilon \to 0^+$ and so $\hat{u}_{\varepsilon}^j \in \theta_{\varepsilon}^j$ by decreasing $\varepsilon^* > 0$ if necessary. So, decreasing $\delta^* > 0$ if necessary, we have that

$$\beta_{\varepsilon}^{J} < \mathcal{I}_{0,a} + \Upsilon, \ \forall \varepsilon \in (0, \varepsilon^{*})$$

which is the first part of the lemma. To reach the remaining one, if $u \in \partial \theta_{\varepsilon}^{j}$, then

$$u \in S(a)$$
 and $|Q_{\varepsilon}(u) - x^j| = \rho_0 > \frac{\rho}{2}$

leading to $Q_{\varepsilon}(u) \notin K_{\frac{\rho_0}{2}}$. Due to Lemma 3.11, we find that

$$I_{\varepsilon}(u) > \mathcal{I}_{0,a} + \frac{\delta^*}{2}$$
, for all $u \in \partial \theta_{\varepsilon}^j$ and $\varepsilon \in (0, \varepsilon^*)$,

and so

$$\tilde{\beta}_{\varepsilon}^{j} = \inf_{u \in \partial \theta_{\varepsilon}^{j}} I_{\varepsilon}(u) \ge \mathcal{I}_{0,a} + \frac{1}{2} \delta^{*}, \ \forall \varepsilon \in (0, \varepsilon^{*}),$$

from where it follows that

$$\beta_{\varepsilon}^{j} < \tilde{\beta}_{\varepsilon}^{j}$$
, for all $\varepsilon \in (0, \varepsilon^{*})$

finishing the proof of this lemma. \Box

Now, we are in a position to investigate the existence of multiple critical points for I_{ε} constrained on S(a).

Proposition 3.13. Let $2 \le p < N$ and $a > \tilde{a} > 0$. Suppose $\varepsilon \in (0, \varepsilon^*)$ to be fixed, decreasing $\varepsilon^* > 0$ if necessary, then $I_{\varepsilon}|_{S(a)}$ has at least l different nontrivial critical points.

Proof. Given a $j \in \{1, \dots, l\}$, we could exploit the Ekeland's variational principle to find a sequence $\{u_n^j\} \subset S(a)$ satisfying

$$I_{\varepsilon}(u_n^J) \to \beta_{\varepsilon}^j$$

and

$$I_{\varepsilon}(v) - I_{\varepsilon}(u_n^j) \ge -\frac{1}{n} \|v - u_n^j\|, \ \forall v \in \theta_{\varepsilon}^j \text{ with } v \neq u_n^j.$$

It follows from Lemma 3.12 that

$$\beta_{\varepsilon}^{j} < \tilde{\beta}_{\varepsilon}^{j}$$
, for all $\varepsilon \in (0, \varepsilon^{*})$.

and thereby $u_n^j \in \theta_{\varepsilon}^j \setminus \partial \theta_{\varepsilon}^j$ for *n* large enough. For all $v \in T_{u_n^i}S(a) = \{w \in X : \int_{\mathbb{R}^N} |u_n^i|^{p-2}u_n^iwdx = 0\}$, there exists a $\zeta > 0$ such that the path $\gamma : (-\zeta, \zeta) \to S(a)$ defined by

Journal of Differential Equations 421 (2025) 1-49

L. Shen and M. Squassina

$$\gamma(t) = \frac{a(u_n^j + tv)}{|u_n^j + tv|_p}$$

which is of class $C^1((-\zeta, \zeta), S(a))$ and satisfies

$$\gamma(t) \in \theta_{\varepsilon}^{j} \setminus \partial \theta_{\varepsilon}^{j}, \ \forall t \in (-\zeta, \zeta), \ \gamma(0) = u_{n}^{j} \text{ and } \gamma'(0) = v.$$

Hence,

$$I_{\varepsilon}(\gamma(t)) - I_{\varepsilon}(u_n^j) \ge -\frac{1}{n} \|\gamma(t) - u_n^j\|, \ \forall t \in (-\zeta, \zeta),$$

and in particular,

$$\begin{split} \frac{I_{\varepsilon}(\gamma(t)) - I_{\varepsilon}(\gamma(0)))}{t} &= \frac{I_{\varepsilon}(\gamma(t)) - I_{\varepsilon}(u_n^j)}{t} \ge -\frac{1}{n} \left\| \frac{\gamma(t) - u_n^j}{t} \right\| \\ &= -\frac{1}{n} \left\| \frac{\gamma(t) - \gamma(0)}{t} \right\|, \ \forall t \in (0, \zeta). \end{split}$$

Since $I_{\varepsilon} \in \mathcal{C}^1(X, \mathbb{R})$, taking the limit of $t \to 0^+$, we get

$$I_{\varepsilon}'(u_n^j)v \ge -\frac{1}{n} \|v\|.$$

Now, we replace v with -v to obtain

$$\sup\{|I_{\varepsilon}'(u_n^j)v|: \|v\|\leq 1\}\leq \frac{1}{n},$$

leading to

$$I_{\varepsilon}(u_n) \to \beta_{\varepsilon}^j$$
 as $n \to +\infty$ and $||I_{\varepsilon}|'_{S(a)}(u_n)||_{X^*} \to 0$ as $n \to +\infty$,

that is, $\{u_n^j\}$ is a $(PS)_{\beta_{\varepsilon}^j}$ for I_{ε} restricted to S(a). Since $\beta_{\varepsilon}^j < \mathcal{I}_{0,a} + \Upsilon$ by Lemma 3.12, then Theorem 3.10 ensures that there is a u^j such that $u_n^j \to u^j$ in X. Thus,

 $u^j \in \theta^j_{\varepsilon}, \ I_{\varepsilon}(u^j) = \beta^j_{\varepsilon} \text{ and } I_{\varepsilon}|'_{S(a)}(u^j) = 0.$

Owing to the following facts

$$Q_{\varepsilon}(u^{i}) \in \overline{B_{\rho_{0}}(x^{i})}, \ Q_{\varepsilon}(u^{j}) \in \overline{B_{\rho_{0}}(x^{j})}$$

and

$$\overline{B_{\rho_0}(x^i)} \cap \overline{B_{\rho_0}(x^j)} = \emptyset \text{ for } i \neq j,$$

we conclude that $u^i \neq u^j$ for $i \neq j$ while $1 \leq i, j \leq l$. Therefore, I_{ε} has at least l nontrivial critical points (u^j, λ^j) with $\lambda^j < 0$ for all $\varepsilon \in (0, \varepsilon^*)$. The proof is completed. \Box

In order to study the concentrating behavior of positive solutions for (1.1)-(1.2), we shall depend on the obtained solutions of Problem (1.10). According to Proposition 3.13, for all $2 \le p < N$ and $a > \tilde{a} > 0$ and decreasing $\varepsilon^* > 0$ if necessary, there are *l* couples of $(v_{\varepsilon}^j, \lambda_{\varepsilon}^j) \in X \times \mathbb{R}$ such that

$$v_{\varepsilon}^{j} \in \theta_{\varepsilon}^{j}, \ I_{\varepsilon}(v_{\varepsilon}^{j}) = \beta_{\varepsilon}^{j} \text{ and } I_{\varepsilon}'(v_{\varepsilon}^{j}) - \lambda_{\varepsilon}^{j}\Psi'(v_{\varepsilon}^{j}) = 0 \text{ in } X^{*},$$

where $j \in \{1, \dots, l\}$, $v_{\varepsilon}^{j}(x) > 0$ for all $x \in \mathbb{R}^{N}$ and $\lambda^{j} < 0$.

Lemma 3.14. Let $2 \le p < N$ and $a > \tilde{a} > 0$. Suppose $\varepsilon \in (0, \varepsilon^*)$ to be fixed, decreasing $\varepsilon^* > 0$ if necessary, there are $y_{\varepsilon}^j \in \mathbb{R}^N$, $R_0^j > 0$ and $\beta_0^j > 0$ such that

$$\int_{B_{R_0}(y_{\varepsilon}^j)} |v_{\varepsilon}^j|^p dx \ge \beta_0^j,$$

for $j \in \{1, \dots, l\}$. Moreover, the family $\{\varepsilon y_{\varepsilon}^{j}\}$ is bounded and, passing to a subsequence if necessary, $\varepsilon y_{\varepsilon}^{j} \to x^{j}$ as $\varepsilon \to 0^{+}$.

Proof. If it is not the case, there is a sequence $\{\varepsilon_n\}$ with $\varepsilon_n \to 0^+$ such that

$$\lim_{n \to \infty} \sup_{y \in \mathbb{R}^N} \int_{B_r(y)} |v_{\varepsilon_n}^j|^p dx = 0$$

for all R > 0. By means of Lion's Vanishing lemma, we would have that $v_{\varepsilon_n}^j \to 0$ in $L^q(\mathbb{R}^N)$ for each $p < q < p^*$ leading to $F_2(u_n) \to 0$ in $L^1(\mathbb{R}^N)$ by property-(\mathbf{P}_2). Owing to $F_1(s) \ge 0$ for all $s \in \mathbb{R}$ by property-(\mathbf{P}_1), there holds $\lim_{n \to \infty} I_{\varepsilon_n}(v_{\varepsilon_n}^j) \ge 0$ which contradicts with the fact that

$$\lim_{n \to \infty} I_{\varepsilon_n}(v_{\varepsilon_n}^j) = \lim_{n \to \infty} \beta_{\varepsilon_n}^i \le \mathcal{I}_{0,a} + \Upsilon < 0.$$
(3.13)

So, we can define $\bar{v}_{\varepsilon}^{j}(\cdot) = v_{\varepsilon}^{j}(\cdot + y_{\varepsilon}^{j})$ and $\{\bar{v}_{\varepsilon}^{j}\}$ is bounded with respect to $\varepsilon \in (0, \varepsilon^{*})$. Therefore, there is a $\bar{v} \in X \setminus \{0\}$ such that $\bar{v}_{\varepsilon}^{j} \rightarrow \bar{v}^{j}$ in X as $\varepsilon \rightarrow 0^{+}$ along a subsequence. Since $\{\bar{v}_{\varepsilon}^{j}\} \subset S(a)$ and

$$I_{\varepsilon}(v_{\varepsilon}^{j}) \geq I_{0}(v_{\varepsilon}^{j}) = I_{0}(\bar{v}_{\varepsilon}^{j}) \geq \mathcal{I}_{0,a}$$

jointly with (3.12) yields that $\lim_{\varepsilon \to 0^+} I_0(\bar{v}^j_\varepsilon) = \mathcal{I}_{0,a}$. Recalling Theorem 3.5, we know that $\bar{v}^j_\varepsilon \to \bar{v}$ in X as $\varepsilon \to 0^+$. Suppose that $\{\varepsilon y^j_\varepsilon\}$ is unbounded with respect to $\varepsilon \in (0, \varepsilon^*)$ and so we can assume that there exists a subsequence $\{\varepsilon_n y^j_{\varepsilon_n}\}$ such that $|\varepsilon_n y^j_{\varepsilon_n}| \to +\infty$ as $n \to \infty$. Exploiting $\bar{v}^j_{\varepsilon_n} \to \bar{v}$ in X,

Journal of Differential Equations 421 (2025) 1-49

$$\begin{split} I_{\varepsilon_n}(v_{\varepsilon_n}^j) &= \frac{1}{p} \int\limits_{\mathbb{R}^N} \left[|\nabla v_{\varepsilon_n}^j|^p + (V(\varepsilon_n x) + 1)|v_{\varepsilon_n}^j|^p \right] dx + \int\limits_{\mathbb{R}^N} F_1(v_{\varepsilon_n}^j) dx - \int\limits_{\mathbb{R}^N} F_2(v_{\varepsilon_n}^j) dx \\ &= \frac{1}{p} \int\limits_{\mathbb{R}^N} \left[|\nabla \bar{v}_{\varepsilon_n}^j|^p + (V(\varepsilon_n x + \varepsilon_n y_n) + 1)|\bar{v}_{\varepsilon_n}^j|^p \right] dx + \int\limits_{\mathbb{R}^N} F_1(\bar{v}_{\varepsilon_n}^j) dx - \int\limits_{\mathbb{R}^N} F_2(\bar{v}_{\varepsilon_n}^j) dx \\ &\to I_{\infty}(\bar{v}) \end{split}$$

together with (3.13) reveals the following inequality

$$\mathcal{I}_{0,a} + \Upsilon \ge I_{\infty}(\bar{v}) \ge \mathcal{I}_{\infty,a}$$

Due to (3.9), it is impossible by the definition of Υ appearing in Theorem 3.10. Therefore, up to a subsequence if necessary, $\varepsilon y_{\varepsilon}^{j} \to x_{0}^{j}$ in \mathbb{R}^{N} as $\varepsilon \to 0^{+}$ and then the remaining part is to verify that $x_{0}^{j} = x^{j}$. Actually, we could use a similar argument, or follow the method adopted in the case ii)-(2) in the proof of Lemma 3.11, to conclude that $V(x_{0}^{j}) = V_{0}$. Recalling $v_{\varepsilon}^{j} \in \theta_{\varepsilon}^{j}$ and it would be simple to see that $\lim_{n \to \infty} Q_{\varepsilon_{n}}(v_{\varepsilon}^{j}) = x_{0}^{j}$, one has that $|x^{j} - x_{0}^{j}| \le \rho_{0}$. Hence, we must have that $x_{0}^{j} = x^{j}$. The proof is completed. \Box

Lemma 3.15. Let $2 \le p < N$ and $a > \tilde{a} > 0$. Suppose $\varepsilon \in (0, \varepsilon^*)$ to be fixed, decreasing $\varepsilon^* > 0$ if necessary, then v_{ε}^j possesses a maximum η_{ε}^j satisfying $V(\varepsilon \eta_{\varepsilon}^j) \to V(x^j)$ as $\varepsilon \to 0^+$ for $j \in \{1, 2, \dots, l\}$. Moreover, there exist $C_0^j, c_0^j > 0$ such that

$$v_{\varepsilon}^{j}(x) \leq C_{0}^{j} \exp(-c_{0}^{j}|x-\eta_{\varepsilon}^{j}|)$$

for all $\varepsilon \in (0, \varepsilon^*)$ and $x \in \mathbb{R}^N$.

Proof. Firstly, we analyze some properties of $\bar{v}_{\varepsilon}^{j}$. Since $\bar{v}_{\varepsilon}^{j}(\cdot) = v_{\varepsilon}^{j}(\cdot + y_{\varepsilon}^{j})$, the definition of v_{ε}^{j} reveals that $(\bar{v}_{\varepsilon}^{j}, \lambda_{\varepsilon}^{j})$ is a couple of weak solution to the problem

$$\begin{cases} -\Delta_p \bar{v}_{\varepsilon}^j + V(\varepsilon x + \varepsilon x_{\varepsilon}^j) |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j = \lambda_{\varepsilon}^j |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j + |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j \log |\bar{v}_{\varepsilon}^j|^p \text{ in } \mathbb{R}^N, \\ \int_{\mathbb{R}^N} |\bar{v}_{\varepsilon}^j|^p dx = a^p. \end{cases}$$
(3.14)

Recalling the arguments explored in Proposition 3.13 and Lemma 3.14, we derive $\bar{v}_{\varepsilon}^{j} \rightarrow \bar{v}^{j}$ in X, $\lambda_{\varepsilon}^{j} \rightarrow \lambda^{j}$ in \mathbb{R} and $\varepsilon x_{\varepsilon}^{j} \rightarrow x^{j}$ in \mathbb{R}^{N} as $\varepsilon \rightarrow 0^{+}$. So, using (3.14), $(\bar{v}^{j}, \lambda^{j})$ is a nontrivial solution to

$$-\Delta_p v + V_0 |v|^{p-2} v = \lambda |v|^{p-2} v + |v|^{p-2} v \log |v|^p \text{ in } \mathbb{R}^N.$$

Similar to Step 1 in the proof of Theorem 2.5, there holds $\bar{v}^j \in L^{\infty}(\mathbb{R}^N)$. Hence, $\bar{v}^j_{\varepsilon} \in L^{\infty}(\mathbb{R}^N)$ and there is a constant C > 0 independent of ε such that $|\bar{v}^j_{\varepsilon}|_{\infty} \leq C$. Indeed, one can further deduce that $\bar{v}^j_{\varepsilon} \in C^{1,\tau}_{\text{loc}}(\mathbb{R}^N)$ for some $\tau \in (0, 1)$. We postpone the detailed proofs in Lemma A.2 in the Appendix to give that

$$|\bar{v}_{\varepsilon}^{j}|_{\infty} \ge \rho^{j}$$
 and $\lim_{|x| \to +\infty} \bar{v}_{\varepsilon}^{j}(x) = 0$ uniformly in $\varepsilon \in (0, \varepsilon^{*})$.

where $\rho^j > 0$ is independent of $\varepsilon \in (0, \varepsilon^*)$.

Secondly, we verify that there exist $\bar{C}_0^j, \bar{c}_0^j > 0$ such that $\bar{v}_{\varepsilon}^j(x) \leq \bar{C}_0^j \exp(-\bar{c}_0^j|x|)$ for all $\varepsilon \in (0, \varepsilon^*)$ and $x \in \mathbb{R}^N$, see Lemma A.3 in the Appendix in detail.

Finally, let $\varrho_{\varepsilon}^{j}$ be a maximum of $\bar{v}_{\varepsilon}^{j}$, we have that $|\bar{v}_{\varepsilon}^{j}(\varrho_{\varepsilon}^{j})|_{\infty} \ge \rho^{j}$. Since $\lim_{|x|\to\infty} \bar{v}_{\varepsilon}^{j}(x) = 0$ uniformly in ε , there exists a $M_{0}^{j} > 0$ independent of ε such that $|\varrho_{\varepsilon}^{j}| \le M_{0}^{j}$. Recalling $\bar{v}_{\varepsilon}^{j}(\cdot) = v_{\varepsilon}^{j}(\cdot + y_{\varepsilon}^{j})$, then $y_{\varepsilon}^{j} + \varrho_{\varepsilon}^{j}$ is a maximum of v_{ε}^{j} . Define $\eta_{\varepsilon}^{j} = y_{\varepsilon}^{j} + \varrho_{\varepsilon}^{j}$, according to Lemma 3.14 and $|\varrho_{\varepsilon}^{j}| \le M_{0}^{j}$, we are derived that $\varepsilon \eta_{\varepsilon}^{j} \to x^{j}$ as $\varepsilon \to 0^{+}$ and hence $V(\varepsilon \eta_{\varepsilon}^{j}) \to V(x^{j})$ by the continuity of V. Moreover, since $\bar{v}_{\varepsilon}^{j}(x) \le \bar{C}_{0}^{j} \exp(-\bar{C}_{0}^{j}|x|)$ for all $x \in \mathbb{R}^{N}$ and $|\varrho_{\varepsilon}^{j}| \le M_{0}^{j}$, there holds

$$v_{\varepsilon}^{j}(x) = \bar{v}_{\varepsilon}^{j}(x - y_{\varepsilon}^{j}) \le \bar{C}_{0}^{j}\exp(-\bar{c}_{0}^{j}|x - y_{\varepsilon}^{j}|) = \bar{C}_{0}^{j}\exp(-\bar{c}_{0}^{j}|x - \eta_{\varepsilon}^{j} + \rho_{\varepsilon}^{j}|) \le C_{0}^{j}\exp(-c_{0}^{j}|x - \eta_{\varepsilon}^{j}|)$$

finishing the proof of this lemma. \Box

Proof of Theorem 1.3. By Proposition 3.13 and Lemma 3.15, we see that Problem (1.10) admits at least *l* different couples of solutions $(v_{\varepsilon}^{j}, \lambda_{\varepsilon}^{j}) \in X \times \mathbb{R}$ with $v_{\varepsilon}^{j}(x) > 0$ for every $x \in \mathbb{R}^{N}$ and $\lambda_{\varepsilon}^{j} < 0$, where $j \in \{1, 2, \dots, l\}$. Moreover, there exist $C_{0}^{j}, c_{0}^{j} > 0$ such that

$$v_{\varepsilon}^{j}(x) \leq C_{0}^{j} \exp(-c_{0}^{j}|x-\eta_{\varepsilon}^{j}|)$$

for all $\varepsilon \in (0, \varepsilon^*)$ and $x \in \mathbb{R}^N$. Let $u_{\varepsilon}^j(\cdot) = v_{\varepsilon}^j(\cdot/\varepsilon)$ and $z_{\varepsilon}^j = \varepsilon \eta_{\varepsilon}^j$ for $j \in \{1, 2, \dots, l\}$, then $(u_{\varepsilon}^j, \lambda_{\varepsilon}^j)$ is the desired solution for $j \in \{1, 2, \dots, l\}$ and Theorem 1.3 is proved. \Box

4. The autonomous problem

In this section, we mainly deal with the existence of normalized solutions for a class of autonomous *p*-Laplacian equations with logarithmic nonlinearities.

4.1. The L^p -subcritical case

In this subsection, to study the Problem (1.14), we need the following minimization problems

$$m(a) = \inf_{u \in S(a)} J(u) \text{ and } m_r(a) = \inf_{u \in S_r(a)} J(u),$$

where $S_r(a) = S(a) \cap X_r$ and the variational functional J is defined by (1.15).

In order to prove Theorem 1.4, we are going to introduce the following lemmas.

Lemma 4.1. Let $2 \le p < N$, then the functional J is coercive and bounded from below on S(a) for all a > 0 and there is an constant $a_* > 0$ such that $m(a) \le 0$ for all $a > a_*$. Moreover, $m(a) = m_r(a)$ for all a > 0.

Proof. Repeating the calculations in the proofs of Lemmas 3.2 and 3.3, we can conclude the first part of this lemma and the details are omitted.

Then, we verify that $m(a) = m_r(a)$. Since $S_r(a) \subset S(a)$, one easily sees that $m(a) \leq m_r(a)$. Thus, we just need to prove that $m(a) \geq m_r(a)$. Suppose that $\{u_n\} \subset S(a)$ is a minimizing sequence with respect to m(a). Denoting u_n^* to be the Schwarz symmetric decreasing rearrangement of u_n , so the Pólya-Szegö's inequality yields that $|\nabla u_n^*|_p \leq |\nabla u_n|_p$. Noting that $|u_n^*|_r = |u_n|_r$ for every $r \in [p, p^*]$, we obtain that $\{u_n^*\} \subset S_r(a)$. Since F_1 and F_2 are nondecreasing in $[0, +\infty]$ by property-(\mathbf{P}_1) and property-(\mathbf{P}_2), then the properties of Schwarz rearrangement (see e.g. [41]) imply that

$$\int_{\mathbb{R}^N} F_1(u_n^*) dx = \int_{\mathbb{R}^N} F_1(u_n) dx, \quad \int_{\mathbb{R}^N} F_2(u_n^*) dx = \int_{\mathbb{R}^N} F_2(u_n) dx,$$

from which, by (1.9), there holds

$$\int_{\mathbb{R}^N} |u_n^*|^p \log |u_n^*|^p dx = \int_{\mathbb{R}^N} |u_n|^p \log |u_n|^p dx.$$

As a consequence,

$$m_r(a) = \inf_{u \in S_r(a)} J(u) \le \inf_{u \in S(c)} J(u) = m(a).$$

The proof is completed. \Box

Lemma 4.2. Let
$$2 \le p < N$$
, then $m(\sqrt[p]{a_1^p + a_2^p}) \le m(a_1) + m(a_2)$ for all $a_1, a_2 > 0$.

Proof. In light of the variational functional *J* is invariant under any translation in \mathbb{R}^N , then, adopting the definition of m(a) and the density of $C_0^{\infty}(\mathbb{R}^N)$ in *X*, we deduce that, for every $\epsilon > 0$, there exist two functions $\psi_1, \psi_2 \in C_0^{\infty}(\mathbb{R}^N)$ with supp $\psi_1 \cap \text{supp } \psi_2 = \emptyset$ and $\psi_1 \in S(a_1)$, $\psi_2 \in S(a_2)$ such that

$$J(\psi_1) \le m(a_1) + \frac{1}{2}\epsilon$$
 and $J(\psi_2) \le m(a_2) + \frac{1}{2}\epsilon$. (4.1)

Without loss of generality, we assume that

dist(supp
$$\psi_1$$
, supp ψ_2) $\ge n$ for some $n \in \mathbb{N}^+$. (4.2)

Now define $\psi \triangleq \psi_1 + \psi_2$, since ψ_1 and ψ_2 have disjoint supports, then $\psi \in S(\sqrt[p]{a_1^p + a_2^p})$ and

$$\begin{cases} \int |\nabla \psi|^p dx = \int |\nabla \psi_1|^p dx + \int |\nabla \psi_2|^p dx, \\ \int \\ \mathbb{R}^N \int |\psi|^s dx = \int \\ \mathbb{R}^N |\psi_1|^s dx + \int |\psi_2|^s dx, \quad \forall s \in [p, p^*], \\ \int \\ \mathbb{R}^N \int |\psi|^p \log |\psi|^p dx = \int \\ \mathbb{R}^N |\psi_1|^p \log |\psi_1|^p dx + \int \\ \mathbb{R}^N \int \\ \mathbb{R}^N |\psi_2|^p \log |\psi_2|^p dx. \end{cases}$$
(4.3)

Hence, for $n \in \mathbb{N}^+$ large enough, we are derived from (4.1), (4.2) and (4.3) that

$$m\left(\sqrt[p]{a_1^p + a_2^p}\right) \le J(\psi) = J(\psi_1) + J(\psi_2) \le m(a_1) + m(a_2) + \epsilon,$$

and the proof is completed. \Box

Lemma 4.3. Let $2 \le p < N$, then the mapping $a \mapsto m(a)$ is continuous on $(a_*, +\infty)$, where $a_* > 0$ comes from Lemma 4.1.

Proof. Given an $a > a_*$, without loss of generality, we let $a_n > a_*$ with $a_n \to a$ as $n \to \infty$. For all $n \in \mathbb{N}$, let $\{u_n\} \subset S(a_n)$ such that $J(u_n) \le m(a_n) + \frac{1}{n}$. Thanks to Lemma 4.1, $\{u_n\}$ is uniformly bounded in X and

$$m(a) \le J\left(\frac{a}{a_n}u_n\right) = J(u_n) + o_n(1) \le m(a_n) + o_n(1).$$

On the other hand, given a minimizing sequence $\{v_n\} \subset S(a)$ for m(a), it holds that

$$m(a_n) \le J\left(\frac{a_n}{a}v_n\right) = J(v_n) + o_n(1) = m(a) + o_n(1).$$

The above two facts reveal the desired result and the proof is completed. \Box

Lemma 4.4. Let $2 \le p < N$ and $a > a_*$. Assume that $\{u_n\} \subset S_r(a)$ is a minimizing sequence of $m_r(a)$ with $u_n \rightharpoonup u$ in X_r as $n \rightarrow \infty$. If $u \ne 0$, then $u_n \rightarrow u$ in X_r as $n \rightarrow \infty$.

Proof. Obviously, one sees that $u_n \to u$ in $L^s(\mathbb{R}^N)$ for all $s \in (p, p^*)$ and $|u|_p \le a$ by Fatou's lemma. Owing to property- (\mathbb{P}_2) , we have that $F_2(u_n) \to F_2(u_n)$ in $L^1(\mathbb{R}^N)$. To exhibit the proof clearly, let us divide the proof into two cases.

Case 1. $u_n \to u$ in $L^p(\mathbb{R}^N)$ along a subsequence as $n \to \infty$. In this case, it holds that $u \in S_r(a)$, then we immediately have that

Journal of Differential Equations 421 (2025) 1-49

$$\begin{split} m_r(a) &\leq J(u) = \frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u|^p + |u|^p) dx + \int_{\mathbb{R}^N} F_1(u) dx - \int_{\mathbb{R}^N} F_2(u) dx - \frac{\mu}{q} \int_{\mathbb{R}^N} |u|^q dx \\ &\leq \liminf_{n \to \infty} \left(\frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u_n|^p + |u_n|^p) dx + \int_{\mathbb{R}^N} F_1(u_n) dx - \int_{\mathbb{R}^N} F_2(u_n) dx - \frac{\mu}{q} \int_{\mathbb{R}^N} |u_n|^q dx \right) \\ &= \liminf_{n \to \infty} J(u_n) = m_r(a) \end{split}$$

which yields that

$$\lim_{n \to \infty} \int_{\mathbb{R}^N} (|\nabla u_n|^p + |u_n|^p) dx = \int_{\mathbb{R}^N} (|\nabla u|^p + |u|^p) dx$$

and

$$\lim_{n\to\infty}\int\limits_{\mathbb{R}^N}F_1(u_n)dx=\int\limits_{\mathbb{R}^N}F_1(u)dx.$$

According to $F_1 \in (\Delta_2)$ by Lemma 2.2, then the above two limits provide us that $u_n \to u$ in X_r as $n \to \infty$. The proof is done in this case.

Case 2. $u_n \neq u$ in $L^p(\mathbb{R}^N)$ as $n \neq \infty$. In this case, denoting $d_n \triangleq |u_n - u|_p$, then, up to a subsequence if necessary,

$$\lim_{n \to \infty} |u_n - u|_p^p = \lim_{n \to \infty} d_n^p \stackrel{\text{def}}{=} d^p > 0.$$

It follows from the Brézis-Lieb lemma that $a^p = \lim_{n \to \infty} (d_n^p + |u|_p^p)$. Combining Lemmas 2.4 and 4.2-4.3,

$$m_r(a) = m_r \left(\lim_{n \to \infty} \sqrt[p]{d_n^p} + |u|_p^p \right) = \lim_{n \to \infty} m_r \left(\sqrt[p]{d_n^p} + |u|_p^p \right)$$

$$\leq \lim_{n \to \infty} m_r(d_n) + m_r(|u|_p) \leq \lim_{n \to \infty} J(u_n - u) + J(u)$$

$$= \lim_{n \to \infty} J(u_n) = m_r(a).$$

Proceeding as the proof in Case 1, we see that $u_n \to u$ in X_r along a subsequence as $n \to \infty$. The proof of this lemma is completed. \Box

Proof of Theorem 1.4. First of all, we know that $m(a) \le 0$ for all $a > a_*$. Then, we shall suppose that $\{u_n\} \subset S_r(a)$ is a minimizing sequence for m(a) by Lemma 4.1. Exploiting Lemma 4.1 again, there exists a $u \in X$ such that $u_n \rightharpoonup u$ in X_r along a subsequence. According to Lemma 4.4, the proof is accomplished if we verify that $u \ne 0$. Finally, we are ready to deduce that $u \ne 0$. Arguing it indirectly, we can assume that $u \equiv 0$. For $m(a) \le 0$, since $F_2(u_n) \rightarrow 0$ in $L^1(\mathbb{R}^N)$ by property-(\mathbf{P}_2) and $u_n \rightarrow u$ in $L^s(\mathbb{R}^N)$ for all $s \in (p, p^*)$, then

ā 8 -

$$0 \ge \lim_{n \to \infty} J(u_n) = \lim_{n \to \infty} \left(\frac{1}{p} \int_{\mathbb{R}^N} (|\nabla u_n|^p + |u_n|^p) dx + \int_{\mathbb{R}^N} F_1(u_n) dx \right) \ge \frac{1}{p} a^p > 0$$

which is impossible. Therefore, we arrive at the desired result $u \neq 0$. The positivity of u is trivial, we omit it here. The proof is completed. \Box

4.2. The L^p -supercritical case

In this subsection, we are going to dispose of the Problem (1.14) with $p + \frac{p^2}{N} < q < p^*$ and $\mu > 0$. As explained in Introduction, we need to study Problem (1.16) and so we define the minimization problem

$$m_{\mathcal{R}}(a) = \inf_{u \in S(a)} J_{\mathcal{R}}(u)$$

The same arguments explored in the proof of Theorem 1.4 guarantee that there is $\underline{a}^* > 0$ independent of \mathcal{R} and μ such that $m_{\mathcal{R}}(a) \leq 0$ for all $a > \underline{a}^*$. We would like to point out that it is possible to find such an $\underline{a}^* > 0$ since $J_{\mathcal{R}}(t\psi) \to -\infty$ as $t \to +\infty$ uniformly in \mathcal{R} and μ .

The next result reveals an important estimate involving the norm in X of the solutions $u_{\mathcal{R}}$ for the Problem (1.16).

Lemma 4.5. Let $2 \le p < N$, $p + \frac{p^2}{N} < q < p^*$ and $\mu > 0$. There exists $\underline{\mu}^* = \underline{\mu}^*(\mathcal{R}) > 0$ such that if $\mu \in (0, \underline{\mu}^*)$, then there is C > 0 independent of \mathcal{R} such that the attained function $u_{\mathcal{R}}$ associated with $m_{\mathcal{R}}(a)$ satisfies $|\nabla u_{\mathcal{R}}|_p \le C$ for all $\mathcal{R} > 0$.

Proof. Arguing as in Lemma 3.2, we see that (1.12) combined with (1.17) and Property-(\mathbf{P}_2) with $\tilde{q} = \bar{q}$ gives

$$J_{\mathcal{R}}(u) \geq \frac{1}{p} \int_{\mathbb{R}^{N}} |\nabla u|^{p} dx - (1 + \mu \mathcal{R}^{q-\bar{q}}) C_{\bar{q}} \mathbb{C}_{N,p,\bar{q}} a^{\bar{q}(1-\beta_{\bar{q}})} \left(\int_{\mathbb{R}^{N}} |\nabla u|^{p} dx \right)^{\frac{\bar{q}\beta_{\bar{q}}}{p}}, \ \forall u \in S(a),$$

where $\bar{q} \in \left(p, p + \frac{p^2}{N}\right)$. Fixing $\underline{\mu}^* = \underline{\mu}^*(\mathcal{R}) = \frac{1}{\mathcal{R}^{p-\bar{q}}}$, then for all $\mu \in (0, \underline{\mu}^*)$, one gets

$$J_{\mathcal{R}}(u) \geq \frac{1}{p} \int_{\mathbb{R}^N} |\nabla u|^p dx - 2C_{\bar{q}} \mathbb{C}_{N,p,\bar{q}} a^{\bar{q}(1-\beta_{\bar{q}})} \left(\int_{\mathbb{R}^N} |\nabla u|^p dx \right)^{\frac{q\nu_q}{p}}, \ \forall u \in S(a).$$

Due to $\beta_{\bar{q}}\bar{q} < p$, exploiting the Young's inequality, there is a constant $C_1 > 0$ independent of \mathcal{R} such that

Ξ.R.-

$$\left(\int\limits_{\mathbb{R}^N} |\nabla u|^p dx\right)^{\frac{q,p_q}{p}} \leq C_1 + \frac{1}{4pC_{\bar{q}}\mathbb{C}_{N,p,\bar{q}}a^{\bar{q}(1-\beta_{\bar{q}})}} \int\limits_{\mathbb{R}^N} |\nabla u|^p dx, \ \forall u \in X.$$

Hence, there is a constant $C_2 > 0$ independent of \mathcal{R} such that

$$|\nabla u|_p^p \le 2p J_{\mathcal{R}}(u) + C_2, \quad \forall a > \underline{a}^*, \ \mu \in (0, \mu^*), \ \mathcal{R} > 0 \text{ and } u \in S(a).$$

Since $J_{\mathcal{R}}(u_{\mathcal{R}}) = m_{\mathcal{R}}(a) \leq 0$ for all $a > \underline{a}^*$, we are able to derive the desired result and so the proof is completed. \Box

Proof of Theorem 1.6. By Corollary 1.7, there are $\underline{a}^* > 0$ (independent of \mathcal{R} and μ) and $\underline{\mu}^*$ such that, for all fixed $a > \underline{a}^*$ and $\mu \in (0, \underline{\mu}^*)$, the couple $(u_{\mathcal{R}}^*, \lambda_{\mathcal{R}}^*) \in S_r(a) \times \mathbb{R}$ is a solution of the problem

$$\begin{cases} -\Delta_p u = \lambda |u|^{p-2} u + |u|^{p-2} u \log |u|^p + \mu f_{\mathcal{R}}(u), \text{ in } \mathbb{R}^N, \\ u(x) > 0 \text{ in } \mathbb{R}^N. \end{cases}$$

Since $\mu \in (0, \mu^*)$, the definition of $f_{\mathcal{R}}$ together with (1.17) leads to

$$0 \le \mu f_{\mathcal{R}}(t) \le t^{q-1}, \ \forall t \ge 0 \text{ and } \mathcal{R} > 0.$$

As a consequence, $\{u_{\mathcal{R}}\}\$ is bounded in $L^{s}(\mathbb{R}^{N})$ for all $\mathcal{R} > 0$ and $s \in (p, p^{*})$ by Lemma 4.5 and $\{\lambda_{\mathcal{R}}\}\$ is bounded for all $\mathcal{R} > 0$. Proceeding as the Step 1 in the proof of Theorem 2.5, there is a constant $M \in (0, +\infty)$ that does not depend upon $\mathcal{R} > 0$ satisfying

$$|u_{\mathcal{R}}|_{\infty} \leq M, \ \forall \mathcal{R} > 0.$$

Let us fix $\mathcal{R} > M$, then we know that the couple $(u_{\mathcal{R}}^*, \lambda_{\mathcal{R}}^*) \in X \times \mathbb{R}$ is weak solution for the Problem (1.14) if $a > \underline{a}^*$ and $\mu \in (0, \underline{\mu}^*)$. This finishes the proof of Theorem 1.6. \Box

5. Final comments

Although all of the main results in this article are derived, as far as we are concerned, there are some other interesting questions worth further exploration.

On the one hand, one may naturally wonder that whether the Problems (1.1)-(1.2) admit a ground state solution. To find the ground state, it suffices to study the existence of ground state solutions for Problem (1.10). We say that $u_0 \in X$ is a ground state solution for Problem (1.10) provided that

$$I'_{\varepsilon}(u_0)|_{S(a)} = 0$$
 and $I_{\varepsilon}(u_0) = \inf\{I_{\varepsilon}(u) : I'_{\varepsilon}(u)|_{S(a)} = 0$ and $u \in S(a)\}$,

where the variational functional $I_{\varepsilon}: X \to \mathbb{R}$ defined by (3.8). Although we cannot give an affirmative answer to the above question at present, it will be exhibited as a theorem below.

Theorem 5.1. Let $2 \le p < N$ and $(\hat{V}_1) - (V_2)$. Suppose additionally that the potential V has no other strict global minimum points than $\{x^1, x^2, \dots, x^l\}$. Then, there exist $\hat{a}^* > 0$ and $\hat{\varepsilon}^* > 0$ such that we can derive at least one ground state solution for (1.1)-(1.2) among the solutions obtained by Theorem 1.3 for all $a > \hat{a}^*$ and $\varepsilon \in (0, \hat{\varepsilon}^*)$.

Remark 5.2. If Theorem 5.1 could be proved successfully, one can observe that the ground state solution inherits the properties of concentrating behavior and exponential decay stated in Theorem 1.3. Moreover, the positive parameters \hat{a}^* is larger than a^* and $\hat{\varepsilon}^*$ is smaller than ε^* , respectively.

On the other hand, it would be interesting to handle the existence of solutions for Problem (1.14) perturbed by a mass-critical nonlinearity. More precisely, let us consider the problem below

$$\begin{cases} -\Delta_p u = \lambda |u|^{p-2} u + |u|^{p-2} u \log |u|^p + \mu |u|^{\bar{p}-2} u \text{ in } \mathbb{R}^N, \\ \int_{\mathbb{R}^N} |u|^p dx = a^p. \end{cases}$$
(5.1)

where $\mu > 0$ is a parameter and $\bar{p} = p + \frac{p^2}{N}$. It seems difficult to construct a nontrivial solution for Problem (5.1) in our *p*-Laplacian setting so far because we require the sufficiently small mass *a* to make sure that the variational functional is coercive and bounded from below on *S*(*a*). Whereas, we cannot prove that $m(a) \le 0$ in this situation and it is the essential difference from the classic *p*-Laplacian problems, namely the logarithmic nonlinearity vanishes in Problem (5.1). We conjecture that (5.1) could be solved if one chooses a suitable work space. Speaking it clearly, motivated by [16], there may exist a work space $Z \subset W^{1,p}(\mathbb{R}^N)$ such that $\int_{\mathbb{R}^N} |u|^p |\log |u|^p | dx < \infty$ for all $u \in Z$ and the imbedding $Z \hookrightarrow L^p(\mathbb{R}^N)$ is compact. If it holds true, one would generalize all the main results in [47] to the *p*-Laplacian settings. What's more, the remained case 1 for the results in this paper would be supplemented.

Acknowledgments

The authors are very grateful to the anonymous referee for his/her valuable comments which improved this manuscript a lot. The first author LJS would like to express his sincere thanks to Professor Claudianor Oliveira Alves for the insightful discussions.

Appendix A. Some technical stuff

Lemma A.1. Let $\Omega \subset \mathbb{R}^N$ be an open set, $\mathcal{L} : \Omega \times \mathbb{R} \times \mathbb{R}^N \to \mathbb{R}$ is a C^1 function and $f \in L^{\infty}_{loc}(\Omega)$. If $\xi \mapsto \mathcal{L}(x, s, \xi)$ is strictly convex for each $(x, s) \in \Omega \times \mathbb{R}$ and $u : \Omega \to \mathbb{R}$ is a locally Lipschitz solution of

$$-div\{\nabla_{\xi}\mathcal{L}(x, u, \nabla u)\} + D_{s}\mathcal{L}(x, u, \nabla u) = f \text{ in } \mathcal{D}'(\Omega).$$

Then

$$\sum_{i,j=1}^{N} \int_{\Omega} D_{i}h_{j} D_{\xi_{i}} \mathcal{L}(x, u, \nabla u) D_{j} u dx - \int_{\mathbb{R}^{N}} \left[(divh) \mathcal{L}(x, u, \nabla u) + h \cdot \nabla_{x} \mathcal{L}(x, u, \nabla u) \right] dx$$

$$= \int_{\mathbb{R}^{N}} (h \cdot \nabla u) f(u) dx, \ \forall h \in \mathcal{C}_{c}^{1}(\Omega, \mathbb{R}^{N}).$$
(A.1)

Lemma A.2. Let $(\bar{v}_{\varepsilon}^{j}, \lambda_{\varepsilon}^{j}) \in X \times \mathbb{R}$ be a couple of weak solution to the Problem (3.14), then

$$|\bar{v}_{\varepsilon}^{j}|_{\infty} \ge \rho^{j} \text{ and } \lim_{|x| \to +\infty} \bar{v}_{\varepsilon}^{j}(x) = 0 \text{ uniformly in } \varepsilon \in (0, \varepsilon^{*}).$$

where $\rho^j > 0$ is independent of $\varepsilon \in (0, \varepsilon^*)$.

Proof. If the first part is false, we suppose that $|\bar{v}_{\varepsilon}^{j}|_{\infty} \to 0$ as $\varepsilon \to 0^{+}$ in the sense of a subsequence. Then, it is simple to verify that $\bar{v}_{\varepsilon}^{j} \to 0$ in X which is absurd and thus we just show the second part in detail. For every R > 0 and $0 < r \leq \frac{R}{2}$, we choose a cutoff function $\eta \in C_{0}^{\infty}(\mathbb{R}^{N}, [0, 1])$ such that $\eta(x) = 1$ if $|x| \ge R$, and $\eta(x) = 0$ if $|x| \le R - r$ as well as $|\nabla \eta| \le \frac{2}{r}$. Given $\varepsilon \in (0, \varepsilon^{*})$ and L > 1, define

$$\bar{v}_{\varepsilon,L}^{j}(x) = \begin{cases} \bar{v}_{\varepsilon}^{j}(x), & \bar{v}_{\varepsilon}^{j}(x) \leq L, \\ L, & \bar{v}_{\varepsilon}^{j}(x) \geq L, \end{cases}$$

and

$$\bar{z}_{\varepsilon,L}^{j} = \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} \bar{v}_{\varepsilon}^{j}$$
 and $\bar{w}_{\varepsilon,L}^{j} = \eta \bar{v}_{\varepsilon}^{j} (\bar{v}_{\varepsilon,L}^{j})^{\vartheta-1}$

with $\vartheta > 1$ to be determined later. Taking $\bar{z}_{\varepsilon,L}^{j}$ as a test function in (3.14), we obtain

$$\begin{split} \int_{\mathbb{R}^{N}} \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} |\nabla \bar{v}_{\varepsilon}^{j}|^{p} dx &= -p(\vartheta-1) \int_{\mathbb{R}^{N}} (\bar{v}_{\varepsilon,L}^{j})^{p\vartheta-p-1} \eta^{p} \bar{v}_{\varepsilon}^{j} |\nabla \bar{v}_{\varepsilon}^{j}|^{p-2} \nabla \bar{v}_{\varepsilon}^{j} \nabla \bar{v}_{\varepsilon,L}^{j} dx \\ &- p \int_{\mathbb{R}^{N}} \eta^{p-1} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} \bar{v}_{\varepsilon}^{j} |\nabla \bar{v}_{\varepsilon}^{j}|^{p-2} \nabla \bar{v}_{\varepsilon}^{j} \nabla \eta dx \\ &+ \int_{\mathbb{R}^{N}} f(\bar{v}_{\varepsilon}^{j}) \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} \bar{v}_{\varepsilon}^{j} dx \\ &- \int_{\mathbb{R}^{N}} V_{\varepsilon}(x) |\bar{v}_{\varepsilon}^{j}|^{p} \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} dx, \end{split}$$

where $V_{\varepsilon}(x) = V(\varepsilon x + \varepsilon x_{\varepsilon}^{j})$ and

$$f(\bar{v}^j_{\varepsilon}) = \lambda^j_{\varepsilon} |\bar{v}^j_{\varepsilon}|^{p-2} \bar{v}^j_{\varepsilon} + |\bar{v}^j_{\varepsilon}|^{p-2} \bar{v}^j_{\varepsilon} \log |\bar{v}^j_{\varepsilon}|^p = (\lambda^j_{\varepsilon} - 1) |\bar{v}^j_{\varepsilon}|^{p-2} \bar{v}^j_{\varepsilon} + F'_2(\bar{v}^j_{\varepsilon}) - F'_1(\bar{v}^j_{\varepsilon}).$$

It follows from property-(\mathbf{P}_1) and property-(\mathbf{P}_2) with $\tilde{q} \in (p, p^*)$ that

$$f(\bar{v}^j_{\varepsilon})\bar{v}^j_{\varepsilon} \le (\lambda^j_{\varepsilon} - 1)|\bar{v}^j_{\varepsilon}|^p + pC_{\tilde{q}}|\bar{v}^j_{\varepsilon}|^{\tilde{q}}$$

which indicates that

$$\int_{\mathbb{R}^{N}} \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} |\nabla \bar{v}_{\varepsilon}^{j}|^{p} dx \leq p \int_{\mathbb{R}^{N}} \eta^{p-1} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} \bar{v}_{\varepsilon}^{j} |\nabla \bar{v}_{\varepsilon}^{j}|^{p-1} |\nabla \eta| dx$$
$$+ p C_{\tilde{q}} \int_{\mathbb{R}^{N}} \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} \bar{v}_{\varepsilon}^{j} |\bar{v}_{\varepsilon}^{j}|^{\tilde{q}} dx + (\lambda_{\varepsilon}^{j} - 1 - V_{0}) \int_{\mathbb{R}^{N}} |\bar{v}_{\varepsilon}^{j}|^{p} \eta^{p} (\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)} dx.$$

Using the Young's inequality, it holds that

$$\begin{split} &\int\limits_{\mathbb{R}^N} \left[\eta^p (\bar{v}^j_{\varepsilon,L})^{p(\vartheta-1)} |\nabla \bar{v}^j_{\varepsilon}|^p - (\lambda^j_{\varepsilon} - 1 - V_0) |\bar{v}^j_{\varepsilon}|^p \eta^p (\bar{v}^j_{\varepsilon,L})^{p(\vartheta-1)} \right] dx \\ &\leq C_p \int\limits_{\mathbb{R}^N} (\bar{v}^j_{\varepsilon,L})^{p(\vartheta-1)} |\bar{v}^j_{\varepsilon}|^p |\nabla \eta|^p dx + p C_p C_{\tilde{q}} \int\limits_{\mathbb{R}^N} \eta^p (\bar{v}^j_{\varepsilon,L})^{p(\vartheta-1)} |\bar{v}^j_{\varepsilon}|^{\tilde{q}} dx \end{split}$$

In view of the proof of Theorem 3.10, we obtain that $\lambda_{\varepsilon}^{j} \leq (\lambda^{*})^{j} < 0$ for all $\varepsilon \in (0, \varepsilon^{*})$. Exploiting (\hat{V}_{1}) , there holds $V_{0} + 1 \geq 0$. Moreover, some simple calculations show that

$$|\nabla \bar{w}_{\varepsilon,L}^{j}|^{p} \leq C_{p}\vartheta^{p}\left(\eta^{p}(\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)}|\nabla \bar{v}_{\varepsilon}^{j}|^{p} + |\nabla \eta|^{p}(\bar{v}_{\varepsilon,L}^{j})^{p(\vartheta-1)}|\bar{v}_{\varepsilon}^{j}|^{p}\right).$$

The above facts together with the Sobolev inequality imply that

$$\begin{split} \left(\int\limits_{\mathbb{R}^N} |\bar{w}_{\varepsilon,L}^j|^{p^*} dx \right)^{\frac{p}{p^*}} &\leq \tilde{C}_p \vartheta^p \left(\int\limits_{\mathbb{R}^N} (\bar{v}_{\varepsilon,L}^j)^{p(\vartheta-1)} |\bar{v}_{\varepsilon}^j|^p |\nabla \eta|^p dx + \int\limits_{\mathbb{R}^N} \eta^p (\bar{v}_{\varepsilon,L}^j)^{p(\vartheta-1)} |\bar{v}_{\varepsilon}^j|^{\tilde{q}} dx \right) \\ &\leq \tilde{C}_{p,r} \vartheta^p \left(\int\limits_{R-r \leq |x| \leq R} |\bar{v}_{\varepsilon}^j|^{p\vartheta} dx + \int\limits_{|x| \geq R-r} (\bar{v}_{\varepsilon}^j)^{p(\vartheta-1)} |\bar{v}_{\varepsilon}^j|^{\tilde{q}} dx \right). \end{split}$$

Hereafter, we shall fix $t = \sqrt{r}$, $p^* > \frac{pt}{t-1}$ and $\chi = \frac{p^*(t-1)}{pt} > 1$. As a consequence,

$$\begin{split} \left(\int\limits_{\mathbb{R}^N} |\bar{w}_{\varepsilon,L}^j|^{p^*} dx \right)^{\frac{p}{p^*}} &\leq \tilde{C}_p \vartheta^p \left\{ \left(\int\limits_{R-r \leq |x| \leq R} |\bar{v}_{\varepsilon}^j|^{\frac{p\vartheta_t}{t-1}} dx \right)^{\frac{t-1}{t}} \left(\int\limits_{R-r \leq |x| \leq R} dx \right)^{\frac{1}{t}} \\ &+ \left(\int\limits_{|x| \geq R-r} |\bar{v}_{\varepsilon}^j|^{\frac{p\vartheta_t}{t-1}} dx \right)^{\frac{t-1}{t}} \left(\int\limits_{|x| \geq R-r} |\bar{v}_{\varepsilon}^j|^{(\tilde{q}-p)t} dx \right)^{\frac{1}{t}} \right\} \end{split}$$

Choosing $\tilde{q} = \frac{p(1+t)}{t}$ and applying the Sobolev inequality, we must have that

$$\left(\int_{\mathbb{R}^N} |\bar{w}_{\varepsilon,L}^j|^{p^*} dx\right)^{\frac{p}{p^*}} \leq \tilde{C}_{p,a} \vartheta^p \left(\int_{|x|\geq R-r} |\bar{v}_{\varepsilon}^j|^{\frac{p\vartheta t}{t-1}} dx\right)^{\frac{t-1}{t}}.$$

From which, since $\bar{w}_{\varepsilon,L}^{j} = \eta \bar{v}_{\varepsilon}^{j} (\bar{v}_{\varepsilon,L}^{j})^{\vartheta-1}$, we can derive that

$$\begin{split} \left(\int\limits_{|x|\geq R} |\bar{v}_{\varepsilon,L}^{j}|^{p^{*}\vartheta} dx \right)^{\frac{p}{p^{*}}} &\leq \left(\int\limits_{|x|\geq R} \eta^{p^{*}} |\bar{v}_{\varepsilon}^{j}|^{p^{*}} |\bar{v}_{\varepsilon,L}^{j}|^{p^{*}(\vartheta-1)} dx \right)^{\frac{p}{p^{*}}} \\ &\leq \left(\int\limits_{\mathbb{R}^{N}} |\bar{w}_{\varepsilon,L}^{j}|^{p^{*}} dx \right)^{\frac{p}{p^{*}}} &\leq \tilde{C}_{p,a} \vartheta^{p} \left(\int\limits_{|x|\geq R-r} |\bar{v}_{\varepsilon}^{j}|^{\frac{p\vartheta_{l}}{l-1}} dx \right)^{\frac{l-1}{l}}. \end{split}$$

Letting $L \to +\infty$ in the above inequality, there holds

$$\left(\int_{|x|\geq R} |\bar{v}_{\varepsilon}^{j}|^{p^{*}\vartheta} dx\right)^{\frac{p}{p^{*}}} \leq \tilde{C}_{p,a}\vartheta^{p} \left(\int_{|x|\geq R-r} |\bar{v}_{\varepsilon}^{j}|^{\frac{p\vartheta t}{t-1}} dx\right)^{\frac{t-1}{t}}.$$

Setting $\chi = \frac{p^*(t-1)}{pt}$ and $s = \frac{pt}{t-1}$, we are derived from the above inequality that

$$|\bar{v}_{\varepsilon}^{j}|_{\chi^{m+1}s(|x|\geqslant R)} \leqslant \tilde{C}_{p,a}^{\sum_{i=1}^{m}\chi^{-i}} \chi^{\sum_{i=1}^{m}i\chi^{-i}} |\bar{v}_{\varepsilon}^{j}|_{p^{*}(|x|\geqslant R-r)}$$

and so

$$|\bar{v}_{\varepsilon}^{j}|_{L^{\infty}(|x|\geqslant R)} \leqslant \tilde{C}_{p,a}^{\sum_{i=1}^{m}\chi^{-i}}\chi^{\sum_{i=1}^{m}i\chi^{-i}}|\bar{v}_{\varepsilon}^{j}|_{p^{*}(|x|\geqslant R-r)}.$$

Since $\bar{v}_{\varepsilon}^{j} \rightarrow \bar{v}^{j}$ in X, the last inequality completes the proof. \Box

Lemma A.3. Let $(\bar{v}_{\varepsilon}^{j}, \lambda_{\varepsilon}^{j}) \in X \times \mathbb{R}$ be a couple of weak solution to the Problem (3.14), then there are $C_{0}^{j}, c_{0}^{j} > 0$ such that

$$\bar{v}_{\varepsilon}^{j} \le C_{0}^{j} \exp(-c_{0}^{j}|x|)$$

for all $\varepsilon \in (0, \varepsilon^*)$ and $x \in \mathbb{R}^N$.

Proof. Since we have derived that $\lambda_{\varepsilon}^{j} \to \lambda^{j}$ as $\varepsilon \to 0^{+}$ and $\bar{v}_{\varepsilon}^{j} > 0$ for every $x \in \mathbb{R}^{N}$, we apply Lemma A.2 to deduce that

$$\lim_{|x|\to+\infty} \frac{\lambda_{\varepsilon}^{j} |\bar{v}_{\varepsilon}^{j}|^{p-2} \bar{v}_{\varepsilon}^{j} + |\bar{v}_{\varepsilon}^{j}|^{p-2} \bar{v}_{\varepsilon}^{j} \log |\bar{v}_{\varepsilon}^{j}|^{p}}{|\bar{v}_{\varepsilon}^{j}|^{p-1}} = -\infty \text{ uniformly in } \varepsilon \in (0, \varepsilon^{*}).$$

So, there is an R > 0 which is independent of $\varepsilon \in (0, \varepsilon^*)$ such that

$$\lambda_{\varepsilon}^{j} |\bar{v}_{\varepsilon}^{j}|^{p-2} \bar{v}_{\varepsilon}^{j} + |\bar{v}_{\varepsilon}^{j}|^{p-2} \bar{v}_{\varepsilon}^{j} \log |\bar{v}_{\varepsilon}^{j}|^{p} \le \frac{V_{0}-2}{2} |\bar{v}_{\varepsilon}^{j}|^{p-1}, \ \forall \varepsilon \in (0, \varepsilon^{*}) \text{ and } |x| \ge R.$$

Denoting the constant $\hat{V}_0 = V_0 + 2 \in [1, +\infty)$, then for all $|x| \ge R$, there holds

$$\begin{split} -\Delta_p \bar{v}_{\varepsilon}^j + \frac{\hat{V}_0}{2} |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j &= \lambda_{\varepsilon}^j |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j + |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j \log |\bar{v}_{\varepsilon}^j|^p \\ &- \left[V(\varepsilon x + \varepsilon x_{\varepsilon}^j) - \frac{V_0 + 2}{2} \right] |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j \\ &\leq \lambda_{\varepsilon}^j |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j + |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j \log |\bar{v}_{\varepsilon}^j|^p - \frac{V_0 - 2}{2} |\bar{v}_{\varepsilon}^j|^{p-2} \bar{v}_{\varepsilon}^j \\ &\leq 0. \end{split}$$

Let $\psi^j(x) = C_0^j \exp(-c_0^j |x|)$ with $C_0^j, c_0^j > 0$ such that $(c_0^j)^p (p-1) < \frac{\hat{v}}{2}$ and $\bar{v}_{\varepsilon}^j(x) \le C_0^j \exp(-c_0^j R)$ for all |x| = R. It follows from some simple calculations that

$$-\Delta_p \psi^j + \frac{\hat{V}}{2} (\psi^j)^{p-1} = (\psi^j)^{p-1} \left(\frac{\hat{V}}{2} - (c_0^j)^p (p-1) + \frac{N-1}{|x|} (c_0^j)^{p-1} \right) > 0, \text{ for all } |x| \ge R.$$

Define $\Sigma = \{|x| \ge R\} \cap \{\bar{v}_{\varepsilon}^{j} > \psi^{j}\}$, adopting the following inequality

$$(|x|^{s-2}x - |y|^{s-2}y) \cdot (x - y) \ge 0$$
 for all $s > 1$ and $x, y \in \mathbb{R}^N$

and choosing $\phi = \max{\{\bar{v}_{\varepsilon}^{j} - \psi^{j}, 0\}} \in W_{0}^{1,p}(\mathbb{R}^{N} \setminus B_{R}) \cap X$ as a test function in

$$-\Delta_p(\bar{v}^j_{\varepsilon} - \psi^j) + \frac{\hat{V}}{2} [(\bar{v}^j_{\varepsilon})^{p-1} - (\psi^j)^{p-1}] \le 0, \text{ for all } |x| \ge R$$

to conclude that

$$0 \ge \int_{\Sigma} (|\nabla \bar{v}_{\varepsilon}^{j}|^{p-2} \nabla \bar{v}_{\varepsilon}^{j} - |\nabla \psi^{j}|^{p-2} \psi) \nabla \phi dx + \frac{\hat{V}}{2} \int_{\Sigma} \left[(\bar{v}_{\varepsilon}^{j})^{p-1} - (\psi^{j})^{p-1}) \right] \phi dx \ge 0.$$

Therefore, the set $\Sigma \equiv \emptyset$. From which, we know that $\bar{v}_{\varepsilon}^{j} \leq \psi^{j}(x)$ for all $|x| \geq R$ and

$$\bar{v}_{\varepsilon}^{j} \leq \psi^{j}(x) = C_{0}^{j} \exp(-c_{0}^{j}|x|) \text{ for all } |x| \geq R.$$

Exploiting Lemma A.2 again, $|\bar{v}_{\varepsilon}^{j}|_{\infty} \leq C$ and so the above inequality holds true for the whole space \mathbb{R}^{N} by increasing C_{0}^{j} to be large. The proof is completed. \Box

Data availability

No data was used for the research described in the article.

References

- M. Agueh, Sharp Gagliardo-Nirenberg inequalities via *p*-Laplacian type equations, NoDEA Nonlinear Differ. Equ. Appl. 15 (2008) 457–472.
- [2] C.O. Alves, On existence of multiple normalized solutions to a class of elliptic problems in whole ℝ^N, Z. Angew. Math. Phys. 73 (3) (2022) 97.
- [3] C.O. Alves, I.S. da Silva, Existence of multiple solutions for a Schrödinger logarithmic equation via Lusternik-Schnirelmann category, Anal. Appl. 21 (6) (2023) 1477–1516.
- [4] C.O. Alves, I.S. da Silva, Existence of a positive solution for a class of Schrödinger logarithmic equations on exterior domains, Z. Angew. Math. Phys. 75 (3) (2024) 77.
- [5] C.O. Alves, C. Ji, Existence and concentration of positive solutions for a logarithmic Schrödinger equation via penalization method, Calc. Var. Partial Differ. Equ. 59 (2020) 21.
- [6] C.O. Alves, C. Ji, Existence of a positive solution for a logarithmic Schrödinger equation with saddle-like potential, Manuscr. Math. 164 (2021) 555–575.
- [7] C.O. Alves, C. Ji, Multi-bump positive solutions for a logarithmic Schrödinger equation with deepening potential well, Sci. China Math. 65 (8) (2022) 1577–1598.
- [8] C.O. Alves, C. Ji, Multiple normalized solutions to a logarithmic Schrödinger equation via Lusternik-Schnirelmann category, J. Geom. Anal. 34 (7) (2024) 198.
- [9] C.O. Alves, L. Shen, On existence of normalized solutions to some classes of elliptic problems with L^2 -supercritical growth, submitted for publication.
- [10] C.O. Alves, N.V. Thin, On existence of multiple normalized solutions to a class of elliptic problems in whole \mathbb{R}^N via Lusternik-Schnirelmann category, SIAM J. Math. Anal. 55 (2) (2023) 1264–1283.
- [11] A. Ambrosetti, M. Badiale, S. Cingolani, Semiclassical states of nonlinear Schrödinger equations, Arch. Ration. Mech. Anal. 140 (1997) 285–300.
- [12] A. Ambrosetti, A. Malchiodi, Concentration phenomena for nonlinear Schrödinger equations: recent results and new perspectives, in: H. Berestycki, M. Bertsch, F.E. Browder, L. Nirenberg, L.A. Peletier, L. Véron (Eds.), Perspectives in Nonlinear Partial Differential Equations, in: Contemp. Math., vol. 446, Amer. Math. Soc., Providence, RI, 2007, pp. 19–30.
- [13] T. Bartsch, N. Soave, A natural constraint approach to normalized solutions of nonlinear Schrödinger equations and systems, J. Funct. Anal. 272 (2017) 4998–5037.
- [14] V. Benci, D. Fortunato, Variational Methods in Nonlinear Field Equations. Solitary Waves, Hylomorphic Solitons and Vortices, Springer Monographs in Mathematics, Springer, Cham, 2014, xviii+250 pp.
- [15] H. Berestycki, T. Cazenave, Instabilité des états stationnaires dans les équations de Schrôdinger et de Klein-Gordon non linéaires, C. R. Acad. Sci., Sér. 1 Math. 293 (1981) 489–492.
- [16] T. Cazenave, Stable solutions of the logarithmic Schrödinger equation, Nonlinear Anal. 7 (1983) 1127–1140.
- [17] T. Cazenave, P.-L. Lions, Orbital stability of standing waves for some nonlinear Schrödinger equations, Commun. Math. Phys. 85 (1982) 549–561.
- [18] P. d'Avenia, E. Montefusco, M. Squassina, On the logarithmic Schrödinger equation, Commun. Contemp. Math. 16 (2) (2014) 1350032.
- [19] P. d'Avenia, M. Squassina, M. Zenari, On fractional logarithmic Schrödinger equations, Math. Methods Appl. Sci. 38 (2015) 5207–5216.
- [20] M. Degiovanni, A. Musesti, M. Squassina, On the regularity of solutions in the Pucci-Serrin identity, Calc. Var. Partial Differ. Equ. 18 (3) (2003) 317–334.
- [21] M. del Pino, J. Dolbeault, The optimal Euclidean L^p-Sobolev logarithmic inequality, J. Funct. Anal. 197 (2003) 151–161.
- [22] M. del Pino, P. Felmer, Multipeak bound states of nonlinear Schrödinger equations, Ann. Inst. Henri Poincaré, Anal. Non Linéaire 15 (1998) 127–149.
- [23] M. del Pino, P. Felmer, Local mountain pass for semilinear elliptic problems in unbounded domains, Calc. Var. Partial Differ. Equ. 4 (1996) 121–137.
- [24] S. Deng, Q. Wu, Normalized solutions for p-Laplacian equation with critical Sobolev exponent and mixed nonlinearities, arXiv:2306.06709.

- [25] S. Deng, Q. Wu, Normalized solutions for *p*-Laplacian equations with potential, arXiv:2310.10510.
- [26] E. DiBenedetto, $C^{1,\gamma}$ local regularity of weak solutions of degenerate elliptic equations, Nonlinear Anal. 7 (1985) 827–850.
- [27] Y.H. Ding, F.H. Lin, Solutions of perturbed Schrödinger equations with critical nonlinearity, Calc. Var. Partial Differ. Equ. 30 (2007) 231–249.
- [28] Y.H. Ding, X.Y. Liu, Semiclassical solutions of Schrödinger equations with magnetic fields and critical nonlinearities, Manuscr. Math. 140 (2013) 51–82.
- [29] J.I. Díaz, Nonlinear Partial Differential Equations and Free Boundaries: I. Elliptic Equations, Research Notes in Mathematics, vol. 106, Pitman, Boston, MA, 1985.
- [30] G. Ferriere, Convergence rate in Wasserstein distance and semiclassical limit for the defocusing logarithmic Schrödinger equation, Anal. PDE 14 (2) (2021) 617–666.
- [31] G. Fibich, The Nonlinear Schrödinger Equation: Singular Solutions and Optical, Collapse Applied Mathematical Sciences, Springer, Berlin, 2015.
- [32] A. Floer, A. Weinstein, Nonspreading wave pachets for the packets for the cubic Schrödinger with a bounded potential, J. Funct. Anal. 69 (1986) 397–408.
- [33] D. Gilbarg, N.S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer, Berlin, 2001.
- [34] N. Ikoma, K. Tanaka, Z.-Q. Wang, C. Zhang, Semi-classical states for logarithmic Schrödinger equations, Nonlinearity 34 (4) (2021) 1900–1942.
- [35] L. Jeanjean, Existence of solutions with prescribed norm for semilinear elliptic equations, Nonlinear Anal. 28 (1997) 1633–1659.
- [36] L. Jeanjean, S. Lu, Nonradial normalized solutions for nonlinear scalar field equations, Nonlinearity 32 (12) (2019) 4942–4966.
- [37] L. Jeanjean, S. Lu, A mass supercritical problem revisited, Calc. Var. Partial Differ. Equ. 59 (5) (2020) 174.
- [38] L. Jeanjean, S. Lu, Normalized solutions with positive energies for a coercive problem and application to the cubicquintic nonlinear Schrödinger equation, Math. Models Methods Appl. Sci. 32 (8) (2022) 1557–1588.
- [39] L. Jeanjean, K. Tanaka, Singularly perturbed elliptic problems with superlinear or asymptotically linear nonlinearities, Calc. Var. Partial Differ. Equ. 21 (2004) 287–318.
- [40] X. Li, Existence of normalized ground states for the Sobolev critical Schrödinger equation with combined nonlinearities, Calc. Var. Partial Differ. Equ. 60 (2021) 1–14.
- [41] E.H. Lieb, M. Loss, Analysis, Graduate Studies in Mathematics, AMS, Providence, Rhode Island, 2001.
- [42] N. Mastorakis, H. Fathabadi, On the solution of *p*-Laplacian for non-Newtonian fluid flow, WSEAS Trans. Math. 8 (2009) 238–245.
- [43] M. Jarosław, J. Schino, Normalized solutions to Schrödinger equations in the strongly sublinear regime, Calc. Var. Partial Differ. Equ. 63 (5) (2024) 137.
- [44] P. Rabinowitz, On a class of nonlinear Schrödinger equations, Z. Angew. Math. Phys. 43 (1992) 270–291.
- [45] M.N. Rao, Z.D. Ren, Theory of Orlicz Spaces, Marcel Dekker, New York, 1985.
- [46] W. Shuai, Two sequences of solutions for the semilinear elliptic equations with logarithmic nonlinearities, J. Differ. Equ. 343 (2023) 263–284.
- [47] W. Shuai, X. Yang, Normalized solutions for logarithmic Schrödinger equation with a perturbation of power law nonlinearity, arXiv:2304.08237.
- [48] N. Soave, Normalized ground states for the NLS equation with combined nonlinearities, J. Differ. Equ. 269 (9) (2020) 6941–6987.
- [49] N. Soave, Normalized ground states for the NLS equation with combined nonlinearities: the Sobolev critical case, J. Funct. Anal. 279 (6) (2020) 108610.
- [50] M. Squassina, A. Szulkin, Multiple solutions to logarithmic Schrödinger equations with periodic potential, Calc. Var. Partial Differ. Equ. 54 (1) (2015) 585–597.
- [51] T. Tao, M. Visan, X. Zhang, The nonlinear Schrödinger equation with combined power-type nonlinearities, Commun. Partial Differ. Equ. 32 (2007) 1281–1343.
- [52] J.-L. Vázquez, A strong maximum principle for some quasilinear elliptic equations, Appl. Math. Optim. 12 (1984) 191–202.
- [53] W. Wang, Q. Li, J. Zhou, Y. Li, Normalized solutions for *p*-Laplacian equations with a L²-supercritical growth, Ann. Funct. Anal. 12 (1) (2021) 9.
- [54] X. Wang, On concentration of positive bound states of nonlinear Schrödinger equations, Commun. Math. Phys. 53 (1993) 229–244.
- [55] Z.-Q. Wang, C.X. Zhang, Convergence from power-law to logarithmic-law in nonlinear scalar field equations, Arch. Ration. Mech. Anal. 231 (2019) 45–61.

- [56] J. Wei, Y. Wu, Normalized solutions for Schrödinger equations with critical Sobolev exponent and mixed nonlinearities, J. Funct. Anal. 283 (2022) 1–46.
- [57] M.I. Weinstein, Nonlinear Schrödinger equations and sharp interpolation estimates, Commun. Math. Phys. 87 (1982/1983) 567–576.
- [58] M. Willem, Minimax Theorems, Birkhäuser Verlag, 1996.
- [59] J. Zhang, C. Lei, J. Lei, The existence and nonexistence of normalized solutions for a *p*-Laplacian equation, Appl. Math. Lett. 148 (2024) 108890.
- [60] Z. Zhang, Variational, Topological and Partial Order Methods with Their Applications, Springer, Berlin, 2013.
- [61] Z.X. Zhang, Z.T. Zhang, Normalized solutions to *p*-Laplacian equations with combined nonlinearities, Nonlinearity 35 (2022) 5621–5663.
- [62] K.G. Zloshchastiev, Logarithmic nonlinearity in the theories of quantum gravity: origin of time and observational consequences, Gravit. Cosmol. 16 (2010) 288–297.