



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Journal of Mathematical Analysis and Applications

journal homepage: www.elsevier.com/locate/jmaa



Boundary behavior for a singular quasi-linear elliptic equation

Marco Squassina

Università degli Studi di Verona, Strada Le Grazie 15, I-37134 Verona, Italy

ARTICLE INFO

Article history:

Received 31 March 2012
 Available online 24 April 2012
 Submitted by V. Radulescu

Keywords:

Singular elliptic equations
 Quasi-linear elliptic equations
 Qualitative behavior

ABSTRACT

We obtain the existence and uniqueness, regularity and boundary behavior for a class of singular quasi-linear elliptic equations in a smooth bounded domain.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction and result

Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain. The aim of this note is to establish the existence, uniqueness and boundary behavior of the solutions to the singular quasi-linear problem

$$\begin{cases} -\operatorname{div}(a(u)Du) + \frac{a'(u)}{2}|Du|^2 = f(u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where $a : (0, +\infty) \rightarrow (0, +\infty)$ is a C^1 function bounded away from zero, $f : (0, +\infty) \rightarrow (0, +\infty)$ is a C^1 function and there exist $a_0 > 0$ and $f_0 > 0$ such that

$$\lim_{s \rightarrow 0^+} a(s)s^{2\mu} = a_0, \quad \text{for some } 0 \leq \mu < 1, \quad (1.2)$$

$$\lim_{s \rightarrow 0^+} f(s)s^\gamma = f_0, \quad \text{for some } \gamma > 1, \quad (1.3)$$

$$2f'(s)a(s) \leq f(s)a'(s), \quad \text{for every } s > 0. \quad (1.4)$$

We refer the reader to [1] and to the references included for an explanation of the interest in and motivations for analyzing these equations. In the one-dimensional case, equations such as (1.1) typically arise in certain problems in fluid mechanics and pseudo-plastic flow (see e.g. [2,3]). In the semi-linear case $a \equiv 1$, various results concerning the existence, uniqueness and asymptotic behavior of the solutions have been obtained in the literature already (see [4–8], the monographs [9,10] and the references therein). Under assumptions (1.2)–(1.4), for the case where $d(x, \partial\Omega)$ denotes the distance of a point x in Ω from the boundary $\partial\Omega$, we shall prove the following:

Theorem 1.1. *Problem (1.1) has a unique solution $u \in C^2(\Omega) \cap C(\bar{\Omega})$ and there exist $\Gamma, \Gamma', \Gamma'' > 0$ with*

$$\Gamma d(x, \partial\Omega)^{\frac{2}{1+\gamma-2\mu}} \leq u(x) \leq \Gamma' d(x, \partial\Omega)^{\frac{2}{1+\gamma-2\mu}}, \quad |Du(x)| \leq \Gamma'' d(x, \partial\Omega)^{\frac{1-\gamma+2\mu}{1+\gamma-2\mu}}, \quad \text{as } d(x, \partial\Omega) \rightarrow 0.$$

Moreover $u \in \operatorname{Lip}(\bar{\Omega})$ if $1 < \gamma \leq 1 + 2\mu$, $u \in C^0, \frac{2}{1+\gamma-2\mu}(\bar{\Omega})$ if $\gamma > 1 + 2\mu$ and $u \in H_0^1(\Omega)$ if $\gamma < 3 - 2\mu$.

E-mail address: marco.squassina@univr.it.

Hence, in some sense, the functions a and f compete for the vanishing rate of the solution and for its gradient upper bound, near the boundary $\partial\Omega$. Furthermore, the range for Lipschitz continuity of u up to the boundary is $\gamma \leq 1 + 2\mu$, $\gamma \neq 1$, and thus enlarged with respect to the one for the semi-linear case, namely $0 < \gamma < 1$; see also Remark 2.2. In [11], the author and Gladiali have jointly recently performed a complete study concerning the existence and qualitative behavior around $\partial\Omega$ of the solutions to the problem

$$\begin{cases} \operatorname{div}(a(u)Du) - \frac{a'(u)}{2}|Du|^2 = f(u) & \text{in } \Omega, \\ u(x) \rightarrow +\infty & \text{as } d(x, \partial\Omega) \rightarrow 0, \end{cases}$$

covering situations where a and f have an exponential, polynomial or logarithmic type growth at infinity and a nonsingular behavior around the origin. In contrast, here we focus on the singular behavior at the origin for a and f with the action of the source f being in some sense predominant at zero as regards the diffusion a , due to the constraint $\gamma > 1 > \mu$. Without loss of generality we assume that a grows as s^k and f decays as s^{-p} as $s \rightarrow +\infty$ for some $k \geq 0$ and $p > 1$, in which case one can also obtain estimates for u and $|Du|$ valid on the whole Ω , as pointed out in Remark 2.2. In the particular case where $a \equiv 1$, problem (1.1) reduces to $-\Delta u = f(u)$ and the above estimates reduce to $\Gamma d(x, \partial\Omega)^{2/(1+\gamma)} \leq u(x) \leq \Gamma' d(x, \partial\Omega)^{2/(1+\gamma)}$ and $|Du(x)| \leq \Gamma'' d(x, \partial\Omega)^{(1-\gamma)/(1+\gamma)}$ as $d(x, \partial\Omega) \rightarrow 0$, consistently with the results of [4,7]. Following the lines of [5,7], some easy adaptations of Theorem 1.1 can be obtained to cover the case of non-autonomous nonlinearities such as $q(x)f(u)$ in place of $f(u)$ and of unbounded domains of \mathbb{R}^N . We leave these further developments to the interested reader. As an example of f and a satisfying (1.2)–(1.4) one can take $f(s) = s^{-\gamma}$, $a(s) = s^{-2\mu}$ for $s \leq s_0$ and $a(s) = \theta(s)$ for $s \geq s_0$ for some $s_0 > 0$, with $\theta \in C^1 \cap L^\infty$ bounded away from zero, $\theta(s_0) = s_0^{-2\mu}$, $\theta'(s_0) = -2\mu s_0^{-2\mu-1}$ and $s\theta'(s) + 2\gamma\theta(s) \geq 0$ for $s \geq s_0$.

2. Proof of the result

In this section, we prove Theorem 1.1. We shall assume that conditions (1.2)–(1.4) hold. In order to get information about the existence, uniqueness and the boundary behavior of the solutions to (1.1), we convert the quasi-linear problem (1.1) into a corresponding semi-linear problem through a change of variable procedure involving the Cauchy problem for $g \in C^2((0, +\infty)) \cap C([0, +\infty))$,

$$\begin{cases} g'(s) = \frac{1}{\sqrt{a(g(s))}}, & \text{for } s > 0, \\ g(0) = 0, \\ g(s) > 0, & \text{for } s > 0. \end{cases} \tag{2.1}$$

Due to the requirement $g > 0$, the solutions of (2.1) are unique and solve $\int_0^{g(s)} \sqrt{a(\xi)} d\xi = s$, for $s > 0$. The solution is global for $s > 0$ since a is bounded away from zero. This procedure was also followed in [11] in the framework of explosive solutions, although there g is C^2 around the origin and defined on \mathbb{R} . Now, since $g \in C^2((0, +\infty)) \cap C([0, +\infty))$ and it is strictly increasing, it is readily seen by a direct computation that $u \in C^2(\Omega) \cap C(\bar{\Omega})$ is a positive solution to (1.1) if and only if $v = g^{-1}(u) \in C^2(\mathcal{D}) \cap C(\bar{\mathcal{D}})$ is a positive solution to $-\Delta v = h(v)$ in Ω , where we have set $h(s) := f(g(s))/\sqrt{a(g(s))}$ for $s > 0$. Let us now obtain the asymptotic behavior of the solution g to problem (2.1) as $s \rightarrow 0^+$ depending of the assigned asymptotic behavior of a as $s \rightarrow 0^+$, given by (1.2). For every $0 \leq \mu < 1$, we have

$$\lim_{s \rightarrow 0^+} \frac{g(s)}{s^{\frac{1}{1-\mu}}} = (1 - \mu)^{\frac{1}{1-\mu}} a_0^{\frac{1}{2(\mu-1)}}. \tag{2.2}$$

In fact, taking into account (2.1), by l'Hôpital's rule we have

$$\begin{aligned} \lim_{s \rightarrow 0^+} \frac{g(s)}{s^{\frac{1}{1-\mu}}} &= (1 - \mu) \lim_{s \rightarrow 0^+} \frac{g'(s)}{s^{\frac{\mu}{1-\mu}}} = (1 - \mu) \lim_{s \rightarrow 0^+} \frac{1}{s^{\frac{\mu}{1-\mu}} \sqrt{a(g(s))}} \\ &= (1 - \mu) \lim_{s \rightarrow 0^+} \frac{g^\mu(s)}{s^{\frac{\mu}{1-\mu}} \sqrt{a(g(s))} g^{2\mu}(s)} = \frac{(1 - \mu)}{\sqrt{a_0}} \left(\lim_{s \rightarrow 0^+} \frac{g(s)}{s^{\frac{1}{1-\mu}}} \right)^\mu, \end{aligned}$$

which yields the claim. Moreover, by virtue of (1.2), (1.3) and (2.2), we have

$$\lim_{s \rightarrow 0^+} \frac{h(s)}{s^{\frac{\mu-\gamma}{1-\mu}}} = \lim_{s \rightarrow 0^+} f(g(s))g(s)^\gamma \frac{1}{\sqrt{a(g(s))}g(s)^{2\mu}} \left[\frac{g(s)}{s^{\frac{1}{1-\mu}}} \right]^{\mu-\gamma} = f_0 a_0^{\frac{1-\gamma}{2(\mu-1)}} (1 - \mu)^{-\frac{\gamma-\mu}{1-\mu}}. \tag{2.3}$$

Observe also that, since $a(s) \sim a_\infty s^k$ as $s \rightarrow +\infty$ and $f(s) \sim f_\infty s^{-p}$ as $s \rightarrow +\infty$ for some $a_\infty, f_\infty > 0$, $k \geq 0$ and $p > 1$, if g still denotes the solution to (2.1), we have three facts:

$$\int_1^{+\infty} \frac{f(g(s))}{\sqrt{a(g(s))}} ds < +\infty, \quad \lim_{s \rightarrow 0^+} \frac{f(g(s))}{\sqrt{a(g(s))}} = +\infty, \quad s \mapsto \frac{f(g(s))}{\sqrt{a(g(s))}} \text{ is nonincreasing.} \tag{2.4}$$

The first property follows immediately from the limit

$$\lim_{s \rightarrow +\infty} \frac{g(s)}{s^{\frac{2}{k+2}}} = \left(\frac{k+2}{2} \frac{1}{\sqrt{a_\infty}} \right)^{\frac{2}{k+2}},$$

which was proved in [11]. The other properties follow by (2.3) and (1.4), respectively. By virtue of (2.2) we now prove that, for every $\gamma > 1$ and $0 \leq \mu < 1$, it holds that

$$\lim_{s \rightarrow 0^+} \frac{\int_{g(s)}^{+\infty} f(\xi) d\xi}{s^{\frac{1-\gamma}{1-\mu}}} = f_0 a_0^{\frac{\gamma-1}{2(1-\mu)}} (\gamma - 1)^{-1} (1 - \mu)^{-\frac{\gamma-1}{1-\mu}}. \tag{2.5}$$

In fact, since $\int_{g(s)}^{+\infty} f(\xi) d\xi \rightarrow +\infty$ for all $\gamma > 1$, it follows that

$$\begin{aligned} \lim_{s \rightarrow 0^+} \frac{\int_{g(s)}^{+\infty} f(\xi) d\xi}{s^{\frac{1-\gamma}{1-\mu}}} &= \frac{\mu - 1}{1 - \gamma} \lim_{s \rightarrow 0^+} \frac{f(g(s))}{s^{\frac{\mu-\gamma}{1-\mu}} \sqrt{a(g(s))}} = \frac{\mu - 1}{1 - \gamma} \lim_{s \rightarrow 0^+} \frac{f(g(s))g^\mu(s)}{s^{\frac{\mu-\gamma}{1-\mu}} \sqrt{a(g(s))g^{2\mu}(s)}} \\ &= \frac{\mu - 1}{(1 - \gamma)\sqrt{a_0}} \lim_{s \rightarrow 0^+} f(g(s))g^\gamma(s) \lim_{s \rightarrow 0^+} \frac{g^{\mu-\gamma}(s)}{s^{\frac{\mu-\gamma}{1-\mu}}} \\ &= \frac{\mu - 1}{(1 - \gamma)\sqrt{a_0}} \frac{f_0}{s^{\frac{\mu-\gamma}{1-\mu}}} \lim_{s \rightarrow 0^+} \frac{g^{\mu-\gamma}(s)}{s^{\frac{\mu-\gamma}{1-\mu}}} = \frac{\mu - 1}{(1 - \gamma)\sqrt{a_0}} \left(\lim_{s \rightarrow 0^+} \frac{g(s)}{s^{\frac{1}{1-\mu}}} \right)^{\mu-\gamma} \\ &= f_0 a_0^{\frac{\gamma-1}{2(1-\mu)}} (\gamma - 1)^{-1} (1 - \mu)^{-\frac{\gamma-1}{1-\mu}}. \end{aligned}$$

Now, for every $\ell \geq 0$, there exists a unique solution $\phi \in C([0, +\infty)) \cap C^2((0, +\infty))$ of the problem

$$\begin{cases} \phi'(s) = \sqrt{\ell^2 + 2 \int_{g(\phi(s))}^{+\infty} f(\xi) d\xi}, & \text{for } s > 0, \\ \phi(0) = 0, \quad \phi(s) > 0, & \text{for } s > 0, \\ \lim_{s \rightarrow +\infty} \phi'(s) = \ell, \quad \lim_{s \rightarrow +\infty} \phi(s) = +\infty. \end{cases} \tag{2.6}$$

To prove this, taking into account (2.4), it is sufficient to apply [7, Lemma 1.3]. Notice that, in particular, the solutions to problem (2.6) locally (namely for every fixed $a > 0$) solve the second-order problem

$$\begin{cases} -\phi''(s) = \frac{f(g(\phi(s)))}{\sqrt{a(g(\phi(s)))}}, & \text{for } 0 < s \leq a, \\ \phi(0) = 0, \quad \phi(s) > 0, & \text{for } 0 < s \leq a. \end{cases} \tag{2.7}$$

We can now prove that, for every $\gamma > 1$, $0 \leq \mu < 1$ and $\ell \geq 0$, it holds that

$$\lim_{s \rightarrow 0^+} \frac{\phi(s)}{s^{\frac{2-2\mu}{1+\gamma-2\mu}}} = \left(\frac{1 + \gamma - 2\mu}{2 - 2\mu} \right)^{\frac{2-2\mu}{1+\gamma-2\mu}} f_0^{\frac{1-\mu}{1+\gamma-2\mu}} a_0^{\frac{\gamma-1}{2(1+\gamma-2\mu)}} (\gamma - 1)^{\frac{\mu-1}{1+\gamma-2\mu}} (1 - \mu)^{\frac{1-\gamma}{1+\gamma-2\mu}}, \tag{2.8}$$

where ϕ denotes the unique solution to (2.6). In fact, by l'Hôpital's rule and (2.5), we obtain

$$\begin{aligned} \lim_{s \rightarrow 0^+} \frac{\phi(s)}{s^{\frac{2-2\mu}{1+\gamma-2\mu}}} &= \lim_{s \rightarrow 0^+} \left[\frac{\phi(s)^{\frac{1+\gamma-2\mu}{2-2\mu}}}{s} \right]^{\frac{2-2\mu}{1+\gamma-2\mu}} = \lim_{s \rightarrow 0^+} \left[\frac{\phi(s)^{\frac{1+\gamma-2\mu}{2-2\mu}}}{\int_0^{\phi(s)} \frac{d\tau}{\sqrt{\ell^2 + \int_{g(\tau)}^{+\infty} f(\xi) d\xi}}} \right]^{\frac{2-2\mu}{1+\gamma-2\mu}} \\ &= \left(\frac{1 + \gamma - 2\mu}{2 - 2\mu} \right)^{\frac{2-2\mu}{1+\gamma-2\mu}} \lim_{s \rightarrow 0^+} \left[\phi(s)^{\frac{\gamma-1}{2-2\mu}} \sqrt{\ell^2 + \int_{g(\phi(s))}^{+\infty} f(\xi) d\xi} \right]^{\frac{2-2\mu}{1+\gamma-2\mu}} \\ &= \left(\frac{1 + \gamma - 2\mu}{2 - 2\mu} \right)^{\frac{2-2\mu}{1+\gamma-2\mu}} \lim_{s \rightarrow 0^+} \left(\frac{\ell^2 + \int_{g(\phi(s))}^{+\infty} f(\xi) d\xi}{\phi(s)^{\frac{1-\gamma}{1-\mu}}} \right)^{\frac{1-\mu}{1+\gamma-2\mu}} \\ &= \left(\frac{1 + \gamma - 2\mu}{2 - 2\mu} \right)^{\frac{2-2\mu}{1+\gamma-2\mu}} \lim_{s \rightarrow 0^+} \left(\frac{\int_{g(s)}^{+\infty} f(\xi) d\xi}{s^{\frac{1-\gamma}{1-\mu}}} \right)^{\frac{1-\mu}{1+\gamma-2\mu}} \\ &= \left(\frac{1 + \gamma - 2\mu}{2 - 2\mu} \right)^{\frac{2-2\mu}{1+\gamma-2\mu}} f_0^{\frac{1-\mu}{1+\gamma-2\mu}} a_0^{\frac{\gamma-1}{2(1+\gamma-2\mu)}} (\gamma - 1)^{\frac{\mu-1}{1+\gamma-2\mu}} (1 - \mu)^{\frac{1-\gamma}{1+\gamma-2\mu}}. \end{aligned}$$

We are now ready to conclude the proof of **Theorem 1.1**. In light of [4, Theorem 1.1], since $h(s) \rightarrow +\infty$ as $s \rightarrow 0^+$ by (2.4) and h is nonincreasing for $s > 0$ by (1.4), there exists a unique positive solution $z \in C^2(\Omega) \cap C(\bar{\Omega})$ to $-\Delta v = h(v)$. Then $g(z) \in C^2(\Omega) \cap C(\bar{\Omega})$ is a positive solution to (1.1). Assume that $u_1, u_2 \in C^2(\Omega) \cap C(\bar{\Omega})$, $u_1, u_2 > 0$ solve (1.1). Then $g^{-1}(u_1), g^{-1}(u_2) > 0$ solve $-\Delta v = h(v)$. By uniqueness, we deduce $g^{-1}(u_1) = g^{-1}(u_2)$, in turn yielding $u_1 = u_2$. By virtue of [4, Theorems 2.2 and 2.5] for any solution v of $-\Delta v = h(v)$ there exist four constants $\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4 > 0$ such that for $d(x, \partial\Omega)$ small enough, $\Lambda_1\phi(d(x, \partial\Omega)) \leq v(x) \leq \Lambda_2\phi(d(x, \partial\Omega))$ and $|Dv(x)| \leq \Lambda_3[d(x, \partial\Omega)h(\Lambda_4\phi(d(x, \partial\Omega))) + \phi(d(x, \partial\Omega))/d(x, \partial\Omega)]$, ϕ being a solution to (2.7). On account of formula (2.8), we can find two constants $\Theta_1, \Theta_2 > 0$ such that, for $d(x, \partial\Omega)$ small enough,

$$\Theta_1 d(x, \partial\Omega)^{\frac{2-2\mu}{1+\gamma-2\mu}} \leq \phi(d(x, \partial\Omega)) \leq \Theta_2 d(x, \partial\Omega)^{\frac{2-2\mu}{1+\gamma-2\mu}}, \tag{2.9}$$

which yield in turn

$$\Lambda_1 \Theta_1 d(x, \partial\Omega)^{\frac{2-2\mu}{1+\gamma-2\mu}} \leq v(x) \leq \Lambda_2 \Theta_2 d(x, \partial\Omega)^{\frac{2-2\mu}{1+\gamma-2\mu}}.$$

Finally, since g is increasing and $u = g(v)$, we have

$$g\left(\Lambda_1 \Theta_1 d(x, \partial\Omega)^{\frac{2-2\mu}{1+\gamma-2\mu}}\right) \leq u(x) \leq g\left(\Lambda_2 \Theta_2 d(x, \partial\Omega)^{\frac{2-2\mu}{1+\gamma-2\mu}}\right).$$

Finally, using (2.2), we obtain the desired controls on u . Now, from (2.9), as $d(x, \partial\Omega)$ is small,

$$\phi(d(x, \partial\Omega))/d(x, \partial\Omega) \leq \Theta_2 d(x, \partial\Omega)^{-\frac{\gamma-1}{1+\gamma-2\mu}}.$$

On account of (2.3) and (2.8), there exists a constant $\Theta_3 > 0$ such that

$$\begin{aligned} & d(x, \partial\Omega)h(\Lambda_4\phi(d(x, \partial\Omega))) \\ &= \Lambda_4^{(\mu-\gamma)/(1-\mu)} d(x, \partial\Omega) \left[\frac{h(\Lambda_4\phi(d(x, \partial\Omega)))}{\Lambda_4^{(\mu-\gamma)/(1-\mu)} \phi^{(\mu-\gamma)/(1-\mu)}(d(x, \partial\Omega))} \right] \phi^{(\mu-\gamma)/(1-\mu)}(d(x, \partial\Omega)) \\ &\leq \Theta_3 d(x, \partial\Omega) d(x, \partial\Omega)^{\frac{2\mu-2\gamma}{1+\gamma-2\mu}} = \Theta_3 d(x, \partial\Omega)^{-\frac{\gamma-1}{1+\gamma-2\mu}}, \end{aligned}$$

for $d(x, \partial\Omega)$ small enough, yielding $|Dv(x)| \leq \Lambda_3(\Theta_2 + \Theta_3)d(x, \partial\Omega)^{-\frac{\gamma-1}{1+\gamma-2\mu}}$. Then, we have

$$\begin{aligned} |Du(x)| &= g'(v(x))|Dv(x)| = \frac{|Dv(x)|}{\sqrt{a(g(v(x)))}} = \frac{|Dv(x)|}{\sqrt{a(g(v(x)))g^{2\mu}(v(x))}} g^\mu(v(x)) \\ &= \frac{1}{\sqrt{a(g(v(x)))g^{2\mu}(v(x))}} \frac{g^\mu(v(x))}{v(x)^{\mu/(1-\mu)}} v(x)^{\mu/(1-\mu)} |Dv(x)| \leq \omega_1 v(x)^{\mu/(1-\mu)} |Dv(x)| \\ &\leq \omega_2 d(x, \partial\Omega)^{\frac{2\mu}{1+\gamma-2\mu}} d(x, \partial\Omega)^{-\frac{\gamma-1}{1+\gamma-2\mu}} = \omega_2 d(x, \partial\Omega)^{\frac{1-\gamma+2\mu}{1+\gamma-2\mu}} \end{aligned}$$

for some $\omega_1, \omega_2 > 0$. In particular, if $1 < \gamma \leq 1 + 2\mu$, it follows that u is Lipschitz continuous up the boundary. If instead $\gamma > 1 + 2\mu$, by the above estimates for u and $|Du|$, we find $\Theta_4 > 0$ such that

$$\begin{aligned} \left| Du^{\frac{1+\gamma-2\mu}{2}}(x) \right| &= \frac{1+\gamma-2\mu}{2} u^{\frac{\gamma-1-2\mu}{2}}(x) |Du(x)| \\ &\leq \Theta_4 d(x, \partial\Omega)^{\frac{\gamma-1-2\mu}{1+\gamma-2\mu}} d(x, \partial\Omega)^{\frac{1-\gamma+2\mu}{1+\gamma-2\mu}} = \Theta_4, \end{aligned}$$

whenever $d(x, \partial\Omega)$ is small enough. In turn, since $u^{\frac{1+\gamma-2\mu}{2}}$ is Lipschitz continuous, $0 < 2/(1 + \gamma - 2\mu) < 1$ and $u = (u^{(1+\gamma-2\mu)/2})^{2/(1+\gamma-2\mu)}$ it follows that u is Hölder continuous up to the boundary $\partial\Omega$ with exponent $2/(1 + \gamma - 2\mu)$, as desired. Finally, concerning the Sobolev regularity of the solution u , observe that in light of [7, Theorem 1.3-J2], a necessary and sufficient condition for v to belong to $H_0^1(\Omega)$ is that

$$\lim_{s \rightarrow 0^+} \int_s^1 \phi(\xi)h(\phi(\xi))d\xi < +\infty,$$

and this, since by (2.3) and (2.8) $\phi(\tau)h(\phi(\tau)) \sim \tau^{\frac{2-2\gamma}{1+\gamma-2\mu}}$ as $\tau \rightarrow 0^+$, is satisfied if and only if $\gamma < 3 - 2\mu$. In turn, $u = g(v) \in H_0^1(\Omega)$ if $\gamma < 3 - 2\mu$ since g is Lipschitz continuous on $(0, +\infty)$, a being bounded away from zero. This concludes the proof of the theorem. \square

Remark 2.1. We know from **Theorem 1.1** that a solution u exists unique and it is Lipschitz continuous up to $\partial\Omega$ provided that $1 < \gamma \leq 1 + 2\mu$. On the other hand, if $\gamma < 1$, combining (2.3) with [4, Theorem 2.25] any solution v of $-\Delta v = h(v)$ is Lipschitz continuous, and so is u , since $u = g(v)$ and g is Lipschitz continuous on $(0, +\infty)$ since a is bounded away from zero.

Remark 2.2. Using [7, Theorem 1.3] in place of [4, Theorems 2.2 and 2.5] we can also state some global estimates for u and $|Du|$ which are valid on the whole $\bar{\Omega}$ and not only in a small neighborhood of the boundary $\partial\Omega$. To be precise, under the assumptions of Theorem 1.1, there exist $\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4 > 0$ with

$$g(\Lambda_1\phi(d(x, \partial\Omega))) \leq u(x) \leq g(\Lambda_2\phi(d(x, \partial\Omega))), \quad \text{for } x \in \bar{\Omega},$$

$$|Du(x)| \leq \Lambda_3 \frac{[d(x, \partial\Omega)h(\Lambda_4\phi(d(x, \partial\Omega))) + \phi(d(x, \partial\Omega))/d(x, \partial\Omega)]}{\sqrt{a(u(x))}}, \quad \text{for } x \in \bar{\Omega},$$

where ϕ denotes the solution to problem (2.6). These formulas are obtained through the monotonicity of g and from $|Du(x)| = |Dv(x)|/\sqrt{a(u(x))}$ for all $x \in \bar{\Omega}$, and follow by the relation $u = g(v)$.

Remark 2.3. Let v be a smooth positive solution to $-\Delta v = h(v)$ in Ω , with $v = 0$ on $\partial\Omega$, where $h > 0$ is as in the proof of Theorem 1.1. Let us consider the map $\psi : [0, +\infty) \rightarrow [0, +\infty)$ defined by

$$\psi(s) := \int_0^s \frac{1}{\sqrt{2\mathcal{H}(\xi)}} d\xi, \quad \mathcal{H}(s) := \int_s^{+\infty} h(\xi) d\xi, \quad \text{for } s > 0.$$

It follows that $\psi'(s) = 1/\sqrt{2\mathcal{H}(s)} > 0$, $\psi''(s) = h(s)/(2\mathcal{H}(s))^{3/2} > 0$ and ψ satisfies $\psi''(s) = (\psi'(s))^3 h(s)$ for all $s > 0$. Put $w := -\psi(v) < 0$; we have $|Dv|^2 = |Dw|^2/(\psi'(v))^2$ as well as $\Delta w = -\psi''(v)|Dv|^2 + \psi'(v)h(v) = \psi'(v)h(v)(1 - |Dw|^2)$, yielding

$$\Delta w = f(w, Dw), \quad f(w, Dw) := -\psi'(\psi^{-1}(-w))h(\psi^{-1}(-w))(1 - |Dw|^2),$$

with $w = 0$ on $\partial\Omega$. Whenever $f > 0$ and $z \mapsto 1/f(z, \cdot)$ is convex, one typically obtains some convexity of w if Ω is convex (see [12]) and in turn some convexity of superlevels of u since $u = g \circ \psi^{-1}(-w)$ and since $s \mapsto g \circ \psi^{-1}(s)$ is strictly increasing. See [13, Section 3] for the particular case $a \equiv 1$ and $f(s) = s^{-\gamma}$.

Acknowledgment

This work was partially supported by PRIN “Variational and Topological Methods in the Study of Nonlinear Phenomena”.

References

[1] M. Squassina, Existence, multiplicity, perturbation, and concentration results for a class of quasi-linear elliptic problems, *Electron. J. Differ. Equ. Monogr.* 7 (2006) 213.
 [2] A. Nachman, A. Callegari, A nonlinear singular boundary value problem in the theory of pseudoplastic fluids, *SIAM J. Appl. Math.* 38 (1980) 275–281.
 [3] C.A. Stuart, Existence theorems for a class of nonlinear integral equations, *Math. Z.* 137 (1974) 49–66.
 [4] M.G. Crandall, P.H. Rabinowitz, L. Tartar, On a Dirichlet problem with a singular nonlinearity, *Comm. Partial Differential Equations* 2 (1977) 193–222.
 [5] A.V. Lair, A.W. Shaker, Classical and weak solutions of a singular semilinear elliptic problem, *J. Math. Anal. Appl.* 211 (1997) 371–385.
 [6] A.C. Lazer, P.J. McKenna, On a singular nonlinear elliptic boundary-value problem, *Proc. Amer. Math. Soc.* 111 (1991) 721–730.
 [7] Zhijun Zhang, Jianguang Cheng, Existence and optimal estimates of solutions for singular nonlinear Dirichlet problems, *Nonlinear Anal.* 57 (2004) 473–484.
 [8] M. Ghergu, V. Radulescu, Multiparameter bifurcation and asymptotics for the singular Lane–Emden–Fowler equation with a convection term, *Proc. Roy. Soc. Edinburgh Sect. A* 135 (2005) 61–84.
 [9] M. Ghergu, V. Radulescu, *Singular Elliptic Problems. Bifurcation and Asymptotic Analysis*, in: Oxford Lecture Series in Mathematics and its Applications, vol. 37, Oxford University Press, 2008, p. 320.
 [10] V. Radulescu, Singular phenomena in nonlinear elliptic problems. From blow-up boundary solutions to equations with singular nonlinearities, in: Michel Chipot (Ed.), *Handbook of Differential Equations: Stationary Partial Differential Equations*, vol. 4, North-Holland Elsevier Science, Amsterdam, 2007, pp. 483–591.
 [11] F. Gladiali, M. Squassina, On explosive solutions for a class of quasi-linear elliptic equations, Preprint.
 [12] B. Kawohl, Rearrangements and Convexity of Level Sets in PDE, in: *Lecture Notes in Mathematics*, vol. 1150, Springer, Berlin, 1985.
 [13] S. Berhanu, F. Gladiali, G. Porru, Qualitative properties of solutions to elliptic singular problems, *J. Inequal. Appl.* 3 (1999) 313–330.